

Kaarina Aarnisalo

Equipment hygiene and risk assessment measures as tools in the prevention of *Listeria monocytogenes* –contamination in food processes



VTT PUBLICATIONS 669

Equipment hygiene and risk assessment measures as tools in the prevention of *Listeria monocytogenes* -contamination in food processes

Kaarina Aarnisalo

Dissertation for the degree of Doctor of Science in Technology to be presented with due permission of the Faculty of Chemistry and Materials Sciences, Helsinki University of Technology for public examination and debate in Auditorium V1 at Helsinki University of Technology (Vuorimiehentie 2, Espoo, Finland) on the 25th of January, 2008, at 12 noon.



ISBN 978-951-38-7069-0 (soft back ed.) ISSN 1235-0621 (soft back ed.)

ISBN 978-951-38-7070-6 (URL: http://www.vtt.fi/publications/index.jsp) ISSN 1455-0849 (URL: http://www.vtt.fi/publications/index.jsp)

Copyright © VTT Technical Research Centre of Finland 2007

JULKAISIJA - UTGIVARE - PUBLISHER

VTT, Vuorimiehentie 3, PL 1000, 02044 VTT puh. vaihde 020 722 111, faksi 020 722 4374

VTT, Bergsmansvägen 3, PB 1000, 02044 VTT tel. växel 020 722 111, fax 020 722 4374

VTT Technical Research Centre of Finland, Vuorimiehentie 3, P.O. Box 1000, FI-02044 VTT, Finland phone internat. +358 20 722 111, fax +358 20 722 4374

VTT, Tietotie 2, PL 1000, 02044 VTT puh. vaihde 020 722 111, faksi 020 722 1000

VTT, Datavägen 2, PB 1000, 02044 VTT tel. växel 020 722 111, fax 020 722 1000

VTT Technical Research Centre of Finland, Tietotie 2, P.O. Box 1000, FI-02044 VTT, Finland phone internat. +358 20 722 111, fax +358 20 722 1000

Technical editing Anni Repo

Edita Prima Oy, Helsinki 2007

Aarnisalo, Kaarina. Equipment hygiene and risk assessment measures as tools in the prevention of *Listeria monocytogenes* -contamination in food processes [Laitehygienia- ja riskinarviointitoimenpiteet työvälineinä *Listeria monocytogenes* -kontaminaation torjumisessa elintarvikeprosesseissa]. Espoo 2007. VTT Publications 669. 101 p. + app. 65 p.

Keywords food hygiene, *Listeria monocytogenes*, risk assessment, hygienic design, disinfectant, lubricant, maintenance, microbial modelling, transfer of bacteria, automated ribotyping

Abstract

Several factors affect on the hygiene level of food processing equipment. A problematic pathogen occurring in food processing is *Listeria monocytogenes*, causing listeriosis with high mortality (20–30%) especially for individuals with reduced immunity. This bacterium is very tolerant to different stress factors and as it can be present in most of the raw materials of food processes, its total elimination is almost impossible. Efficient control of *L. monocytogenes* at the processing plant level requires good equipment hygiene, including functioning good manufacturing and hygiene practices used by all employees, effective means of decontamination and rapid detection of contamination sources, as well as hazard analysis systems supported by risk assessment procedures. The present thesis focuses on deficiencies and improvements in these equipment hygiene and risk assessment practices with the aim of elucidating and developing the most efficient practices against *L. monocytogenes*.

The hygienically most problematic types of equipment in the Finnish food industry were investigated by using a mail-survey. These were identified as the packaging machines, conveyers, dispensers, slicing machines and cooling machines. The main reason for the equipment being considered as problematic was poor hygienic design. The results show clearly that equipment designers must focus their performance on more suitable equipment design. Additionally, an investigation based on a mail-survey and microbiological sampling was made concerning hygiene performance of the maintenance personnel in food processing plants. Clear deficiencies were found e.g. in use of protective clothing, washing of hands and tools as well as avoiding foreign bodies left on the production lines. The results of these studies also indicate that *L. monocytogenes* may be transferred through maintenance work. Training of maintenance personnel with reference to hygienic practices must be increased.

Topics connected to the maintenance operations which have received only minor attention in previous studies include the growth and survival of *L. monocytogenes*

in lubricants used in the equipment as well as control of the bacterium with disinfectants at cold temperatures. In the current thesis the survival, growth and transfer of the bacterium in lubricants used in food processing equipment was studied. The results showed that lubricants used in maintaining the equipment may act as contamination vehicles of *L. monocytogenes*. As the temperatures in food processing premises are usually low, an investigation of the efficiency of eight commonly used commercial disinfectants against *L. monocytogenes* strains at +5 °C was performed. The tested agents were generally efficient at the recommended concentrations and effect times. Thus they appear to be suitable for control of *L. monocytogenes* at the plant level, with only a few exceptions.

Rapid, reliable and easy-to-use methods are needed at the processing plant level. Consequently the suitability of automated ribotyping was compared with the traditionally accepted and successfully used pulsed-field gel electrophoresis (PFGE) to discriminate *L. monocytogenes* isolates and thus trace contamination sources in food plants. PFGE had a higher discriminatory power for *L. monocytogenes* isolates than automated ribotyping. However, based on its automation and rapidity automated ribotyping can be considered a good method for control purposes, although in epidemiological studies identical results must be confirmed with PFGE.

Additionally, in this thesis risk assessment practices were developed by investigating and modelling recontamination of a product and by a plant-level quantitative risk assessment. Transfer of L. monocytogenes from slicing blade to slices of cold-salted salmon was investigated and modelled. Transfer with a progressive exponential reduction in the quantity of bacteria (log CFU/g) in slices was detected. The results provide an example to food processors of how limited data from microbiological analysis can be used to assess the level of recontamination for risk assessment purposes. The principles of microbiological risk assessment can be used at the processing plant level to assist in developing Hazard Analysis Critical Control Point (HACCP)- systems in order to provide a more scientific and comprehensive approach to the control of L. monocytogenes and other microbiological hazards. As a concluding example, a practical approach to quantitative risk assessment of L. monocytogenes for one product at the plant level is presented. This approach helps food processors in illustrating the risks caused by the products for consumers and thus rationalizing risk management actions against L. monocytogenes.

Aarnisalo, Kaarina. Equipment hygiene and risk assessment measures as tools in the prevention of *Listeria monocytogenes* -contamination in food processes [Laitehygienia- ja riskinarviointitoimenpiteet työvälineinä *Listeria monocytogenes* -kontaminaation torjumisessa elintarvikeprosesseissa]. Espoo 2007. VTT Publications 669. 101 s. + liitt. 65 s.

Avainsanat food hygiene, *Listeria monocytogenes*, risk assessment, hygienic design, disinfectant, lubricant, maintenance, microbial modelling, transfer of bacteria, automated ribotyping

Tiivistelmä

Elintarviketuotantolaitteiden hygieniatasoon vaikuttavat useat tekijät. *Listeria monocytogenes* -bakteeri on elintarvikeprosesseissa esiintyvä ongelmallinen patogeeni, joka aiheuttaa listerioosia erityisesti riskiryhmiin kuuluville henkilöille. Tautiin sairastuneiden kuolleisuus on 20–30 %. Bakteeri kestää hyvin erilaisia stressitekijöitä ja koska bakteeria on voitu todeta myös useimmista elintarvike-prosesseissa käytetyistä raaka-aineista, sen täydellinen hävittäminen prosesseista on lähes mahdotonta. *L. monocytogenes* -bakteerin tehokas hallinta tuotantolaitostasolla koostuu hyvästä laitehygieniasta ja sisältää toimivat, hyvät tuotanto- ja hygieniakäytännöt kaikkien työntekijöiden osalta, tehokkaat bakteerin tuhoamiseen tarkoitetut menetelmät ja nopeat kontaminaatiolähteiden toteamismenetelmät yhdistettynä riskinarviointikäytännöillä tuettuun prosessin vaara-analyysiin. Tässä työssä selvitettiin puutteita ja parannuksia nykyisin käytössä olevien laitehygienia-käytäntöjen osalta ja kehitettiin riskinarviointikäytäntöjä mahdollisimman tehok-kaiden *L. monocytogenes* -bakteerin torjumiskeinojen varmistamiseksi.

Tuotantohygienialtaan ongelmallisimpia laitteita Suomen elintarviketeollisuudessa selvitettiin kyselytutkimuksena. Ongelmallisimmiksi laitteiksi todettiin pakkauskoneet, kuljettimet ja annostelu-, siivutus- ja jäähdytyskoneet. Pääasialliseksi syyksi kaikkien näiden laitteiden osalta todettiin huono hygieeninen laitesuunnittelu. Tulokset osoittavat selvästi, että laitevalmistajien on paneuduttava asiaan paremmin ja kehitettävä elintarviketeollisuuteen tarkoitettujen laitteiden hygieenisyyttä. Selvitys tehtiin myös laitteita huoltavan kunnossapitohenkilöstön hygieenisistä toimintatavoista kyselytutkimuksena ja mikrobiologisella näytteenotolla. Selkeitä puutteita toimintatavoissa voitiin todeta mm. suojavaatetuksessa, käsien ja työkalujen puhdistamisessa sekä vierasesineiden tuotantolinjoille jäämisen osalta. Tutkimuksessa löydettiin myös viite siitä, että *L. monocytogenes* voisi siirtyä huoltotyön yhteydessä tehtaan sisällä. Kunnossapitohenkilöstön hygienia-koulutuksen määrää on lisättävä. Kunnossapitotyöhön liittyviä, aiemmin vain vähän tutkittuja aihealueita ovat *L. monocytogenes* -bakteerin kasvu ja selviytyminen laitteissa käytetyissä voiteluaineissa sekä bakteerin tuhoaminen desinfiointiaineilla kylmissä lämpötiloissa. Työssä selvitettiin *L. monocytogenes* -bakteerin selviytymistä, kasvua ja siirtymistä laitteistoissa käytetyissä voiteluaineissa. Tulokset osoittivat, että voiteluaineet voivat toimia *L. monocytogenes* -kontaminaation lähteinä ja siirtymisvälineenä. Koska tuotantolämpötilat elintarvikeprosesseissa ovat yleensä alhaiset, kahdeksan nykyisin käytössä olevan kaupallisen desinfiointiaineen tehoa *L. monocytogenes* -bakteeriin selvitettiin +5 asteessa. Desinfiointiaineet tehosivat yleensä valmistajan antamilla alhaisimmilla käyttökonsentraatioilla ja käyttöajoilla, vain joitakin poikkeuksia lukuun ottamatta.

Koska tehdastasolla tarvitaan nopeita, luotettavia ja helppokäyttöisiä menetelmiä, automaattisen ribotyypityksen soveltuvuutta *L. monocytogenes* -isolaattien erotteluun ja siten kontaminaatiolähteiden jäljittämiseen elintarvikeprosesseissa verrattiin hyväksi todettuun pulssikenttägeelielektroforeesimenetelmään. Tulokset osoittivat, että pulssikenttägeelielektroforeesi oli erottelukyvyltään parempi. Nopeutensa ja automaattisuutensa takia automaattinen ribotyypitys soveltuu kuitenkin myös hyvin hygieniavalvontaan, mutta epidemiologisissa selvityksissä ribotyypityksellä saadut identtiset tulokset on varmistettava pulssikenttägeelielektroforeesilla.

Työssä kehitettiin lisäksi riskinarviointikäytäntöjä selvittämällä ja mallintamalla tuotteen jälkikontaminaatiota sekä kuvaamalla tehdastason riskinarviointi. *L. monocytogenes* -bakteerin siirtymistä siivutettaessa terästä graavilohisiivuihin tutkittiin ja sitä mallinnettiin. Selkeää siirtymistä todettiin bakteerimäärien eksponentiaalisesti pienentyessä siivuissa. Tulokset toimivat esimerkkinä elintarvikeyrityksille rajatun mikrobiologisen tutkimusaineiston käytöstä jälkikontaminaation vaikutuksen arvioimiseen. Kvantitatiivisen mikrobiologisen riskinarvioinnin periaatteita voidaan soveltaa tuotantolaitostasolla Hazard Analysis Critical Control Point (HACCP) -järjestelmien kehittämisessä tieteellisemmän ja kokonaisvaltaisemman listeriavalvonnan aikaansaamiseksi. Tässä työssä viimeisenä esimerkkinä kuvataan käytännönläheinen yhden tuotteen tuotantolinja -tason *L. monocytogenes* -riskinarviointi. Tämä lähestymistapa auttaa elintarviketuottajia hahmottamaan kuluttajille tuotteista aiheutuvaa riskiä ja siten vertailemaan ja rationalisoimaan eri riskinhallintakeinoja *L. monocytogenes* -bakteerin torjumisessa.

Preface

This study was carried out at VTT Technical Research Centre of Finland during the years 2000–2007. The financial support of VTT, Tekes – the Finnish Funding Agency for Technology and Innovation as well as the Jenny and Antti Wihuri Foundation is gratefully acknowledged. All studies were undertaken in cooperative projects with industrial partners, who are also gratefully acknowledged.

I thank Professor Juha Ahvenainen for providing excellent working facilities and Professor Katrina Nordström from the Helsinki University of Technology for her support and cooperation during this study. I am grateful to Professor Raivo Vokk from the Tallinn University of Technology and Dr. Marjatta Rahkio from the Finnish Meat Research Institute for reviewing the thesis manuscript and for their valuable comments.

I am deeply grateful to my supervisor, Technology Manager Dr. Laura Raaska, for her support and encouraging attitude as well as for the time and effort she put into this work. I express my gratitude to my supervisor Dr. Gun Wirtanen for her most valuable advice and comments, especially during the writing process.

My special thanks go to all my co-authors. Dr. Shiowshuh Sheen and Dr. Mark Tamplin are thanked for their excellent guidance and hospitality during my research visit at the United States Department of Agriculture (USDA) in 2006. Dr. Maija-Liisa Suihko is gratefully acknowledged for all the work related to automated ribotyping. My sincere thanks are due to Dr. Riitta Maijala for all her valuable comments, especially those related to risk assessment, as well as to Dr. Jukka Ranta for excellent modelling. Professor Hannu Korkeala is appreciated for his useful comments and pleasant cooperation during the work. I express my gratitude to Dr. Janne Lundén, Dr. Tiina Autio, Elina Vihavainen, DVM, Leila Rantala, M.Sc. (Tech.), Professor Anna-Maija Sjöberg and Dr. Sebastian Hielm for their valuable input to the papers. I also express my special thanks to Pirkko Tuominen, DVM, for her efforts and pleasant cooperation over the years. My colleague and room-mate Dr. Hanna Miettinen and colleagues Dr. Satu Salo, Kirsi Kujanpää, B.Sc., Dr. Hanna-Leena Alakomi, Dr. Outi Priha and Eija Skyttä M.Sc. (Tech.) are thanked for all the fruitful discussions, help which has always been available and their friendship over the years.

Erja Järvinen, Tarja Vappula, Taina Holm, Helena Hakuli, Tarja Niiranen-Jaatinen, Raija Ahonen and Oili Lappalainen are gratefully acknowledged for their invaluable technical assistance. Anne Arvola, M.Sc., is thanked for good advice in statistics and Michael Bailey for revising the English language. I thank other colleagues at VTT for the pleasant work environment. I am also very grateful to all my colleagues at the USDA for all their help and hospitality during my research visit.

I thank all my wonderful friends and relatives for their support and encouragement during the years. Especially, I thank my parents, my mother Leena, who has always been there when needed, and my "special friend" father Timo, for all their support over the years. Special thanks are due to my brother Antti and his family (Piia, Katariina and my god-daughter Elina) and to my brother Pekka with Satu, for their interest on my studies. Finally, I owe my heartfelt thanks to Juha for his ongoing support and patience during this work and for sharing life with me during recent years.

In Espoo, December 2007

Kaarina Aarnisalo

List of publications

- I Aarnisalo, K., Tallavaara, K., Wirtanen, G., Maijala, R. and Raaska, L. 2006. The hygienic working practices of maintenance personnel and equipment hygiene in the Finnish food industry. Food Control, Vol. 17, pp. 1001–1011.
- II Aarnisalo, K., Autio, T., Sjöberg, A.-M., Lundén, J., Korkeala, H. and Suihko, M.-L. 2003. Typing of *Listeria monocytogenes* isolates originating from the food processing industry with automated ribotyping and pulsedfield gel electrophoresis. Journal of Food Protection, Vol. 66, No. 2, pp. 249–255.
- III Aarnisalo, K., Raaska, L. and Wirtanen, G. 2007. Survival and growth of *Listeria monocytogenes* in lubricants used in the food industry. Food Control, Vol. 18, pp. 1019–1025.
- IV Aarnisalo, K., Lundén, J., Korkeala, H. and Wirtanen, G. 2007. Susceptibility of *Listeria monocytogenes* strains to disinfectants and chlorinated alkaline cleaners at cold temperatures. LWT – Food Science and Technology, Vol. 40, pp. 1041–1048.
- V Aarnisalo, K., Sheen, S., Raaska, L. and Tamplin, M. 2007. Modelling transfer of *Listeria monocytogenes* during slicing of 'gravad' salmon. International Journal of Food Microbiology, Vol. 118, pp. 69–78.
- VI Aarnisalo, K., Vihavainen, E., Rantala, L., Maijala, R., Suihko, M.-L., Hielm, S., Tuominen, P., Ranta, J. and Raaska, L. Use of results of microbiological analyses for risk-based control of *Listeria monocytogenes* in marinated broiler legs. International Journal of Food Microbiology, *in press*.

The author's contribution to the appended publications

- I The author had the main responsibility in planning the work on hygienic working practices of maintenance personnel, together with Dr. Laura Raaska; in collecting and analyzing the data and in writing the paper. The author supervised and participated in analyzing the microbiological samples. The author had the main responsibility in planning the questionnaire on hygienic equipment design and in collecting and analyzing the data.
- II The work was planned together with other authors. Kaarina Aarnisalo had the main responsibility in choosing the *L. monocytogenes* isolates for the study and had participated in collecting most of them. She carried out the serotyping, collected the result tables to the paper and compared the results by calculating the *DI* values. She wrote the paper together with Dr. Maija-Liisa Suihko and other authors. Dr. Maija-Liisa Suihko was responsible for the results of automated ribotyping and Dr. Tiina Autio for pulsed-field gel electrophoresis.
- III The author had the main responsibility in designing the experiments, together with Dr. Gun Wirtanen; in interpreting the results and writing the paper. She supervised and partially participated in the laboratory work. Statistical analysis was performed together with Anne Arvola.
- IV The author had the main responsibility in designing the experiments, together with other authors, and in writing the paper. She carried out the transfer experiments. She participated in the modelling work, the main responsibility for modelling being with Dr. Shiowshu Sheen.
- V The author had the main responsibility in planning the work and interpreting the results concerning suspension and surface experiments and in writing the paper. She supervised and partially participated in the laboratory work concerning suspension and surface experiments. Dr. Janne Lundén had main responsibility in the MIC and adaptation experiments.

VI The author had the main responsibility in planning the work and in writing the paper, together with Dr. Laura Raaska and other authors. The author acted as supervisor of and participated in collecting and analyzing the broiler legs and in performing the laboratory heating experiments. The risk assessment was performed together with Dr. Riitta Maijala and Dr. Jukka Ranta (responsible for modelling). Dr. Maija-Liisa Suihko was responsible for the results of automated ribotyping and Leila Rantala for those of pulsed-field gel electrophoresis and serotyping.

Contents

Ab	stract			3
Tii	vistel	mä		5
Pre	eface .			7
Lis	t of p	ublication	ons	9
Th	e auth	or's cor	ntribution to the appended publications	. 10
Lis	t of sy	ymbols.		. 15
1.	Intro	duction		. 17
	1.1	Equipr	nent hygiene in food processing	. 17
		1.1.1	Hygienic design of food processing equipment	.17
		1.1.2	Hygiene of personnel working with equipment	. 18
		1.1.3	Lubricants used in equipment	. 22
		1.1.4	Listeria monocytogenes in equipment hygiene	. 23
	1.2	Legisla	ation and standards	. 23
		1.2.1	Legislation on hygienic design of food processing equipment	23
		1.2.2	Legislation on lubricants used in food processing	. 25
		1.2.3	Legislation on L. monocytogenes	. 26
	1.3	Charac	cteristics of <i>L. monocytogenes</i> – a problematic food	
		proces	sing contaminant	. 26
		1.3.1	The genus Listeria and L. monocytogenes	. 26
		1.3.2	Occurrence	. 27
		1.3.3	Tolerance of pH, temperature and a _w	. 28
		1.3.4	Listeriosis	. 31
		1.3.5	Detection, identification and typing	. 32
		1.3.6	Attachment and transfer	.35
	1.4	Preven	tion of L. monocytogenes in food processes	. 37
		1.4.1	General aspects in prevention	. 37
		1.4.2	Efficiency of disinfectants on L. monocytogenes	. 38
	1.5	Microl	piological risk assessment (MRA) in the food industry	. 42
		1.5.1	Background and components of MRA	. 42

		1.5.2	Microbiological risk assessment of L. monocytogenes	43		
		1.5.3	Use of predictive models in MRA and risk management	43		
		1.5.4	Use of MRA and predictive models at the food plant level.	44		
2.	Aim	s of the	study	46		
3.	Mate	erials ar	nd methods	47		
	3.1	Quest	ionnaires on equipment hygiene and hygienic working			
		practio	ces of maintenance personnel (Paper I)	47		
	3.2	Sampl	ling, detection and identification of L. monocytogenes	48		
		3.2.1	L. monocytogenes strains used in laboratory experiments	48		
		3.2.2	Analyses of tools, work environment and protective			
			clothing of maintenance personnel (Paper I)	49		
		3.2.3	Analyses of cold-salted salmon slices in a transfer study			
			(Paper V)	49		
		3.2.4	Analyses of broiler legs in a risk assessment study (Paper VI).	50		
		3.2.5	Typing of L. monocytogenes (Paper II)	51		
	3.3 Analysing survival and transfer of <i>L. monocytogenes</i> in					
		lubric	ants (Paper III)	51		
		3.3.1	Survival of L. monocytogenes in lubricants	51		
		3.3.2	Transfer from lubricants to stainless steel surfaces			
			and vice versa	52		
	3.4	Susce	ptibility of L. monocytogenes to disinfectants (Paper IV)	54		
		3.4.1	Suspension method	54		
		3.4.2	Surface method	54		
	3.5	Analy	sing transfer of L. monocytogenes during slicing of			
		cold-s	alted salmon (Paper V)	55		
	3.6	Mathe	ematical methods	56		
		3.6.1	Microbicidal effect (Papers III, IV)	56		
		3.6.2	Discrimination index (Paper II)	56		
		3.6.3	Statistical analyses (Papers I, III, V)	57		
		3.6.4	Predictive modelling (Paper V)	57		
		3.6.5	Producer level quantitative risk assessment (Paper VI)	58		
4.	Resi	ults and	discussion	60		
	4.1	Hygie	nically most problematic food processing equipment (Paper 1	I) 60		
	4.2	Hygie	nic working practices of maintenance personnel (Paper I)	62		
		4.2.1	Significant aspects according to the questionnaire	62		

		4.2.2	Microbiological sampling in food processing	. 64		
4.3 Discriminatory power of automated ribotyping compared with						
		PFGE	in distinguishing L. monocytogenes isolates (Paper II)	. 65		
	4.4	Surviv	al and transfer of <i>L. monocytogenes</i> in lubricants	. 66		
		4.4.1	Hygiene of lubricants according to the questionnaire (Paper I)	66		
		4.4.2	Survival of L. monocytogenes in lubricants (Paper III)	. 67		
		4.4.3	Transfer of <i>L. monocytogenes</i> from lubricants to stainless			
			steel surfaces and vice versa (Paper III)	. 68		
	4.5	Suscep	tibility of L. monocytogenes to disinfectants (Paper IV)	. 69		
		4.5.1	Efficacy of disinfectants in suspension	. 69		
		4.5.2	Efficacy of disinfectants on surfaces	. 70		
	4.6	Transf	er of L. monocytogenes during slicing of cold-salted salmon			
		(Paper	V)	. 71		
		4.6.1	Transfer of L. monocytogenes from slicing blade to slices	. 71		
		4.6.2	Model of transfer of <i>L. monocytogenes</i>	. 73		
	4.7	Produc	er level quantitative risk assessment of L. monocytogenes			
		(Paper	VI)	. 74		
		4.7.1	Prevalence and numbers of L. monocytogenes in broiler legs	74		
		4.7.2	Thermal inactivation of L. monocytogenes by heating	. 75		
		4.7.3	Quantitative risk assessment	. 76		
5.	Conc	lusions		. 80		
Re	ferenc	es		. 84		
Ap	pendi	ces				

Papers I-VI

Appendices of this publication are not included in the PDF version. Please order the printed version to get the complete publication (http://www.vtt.fi/publications/index.jsp)

List of symbols

AISI	American Iron and Steel Institute
ALOA	Agar Listeria Ottavani and Agosti
ANSI	American National Standards Institute
a _w	water activity
CCD	charge-coupled device camera
ССР	Critical Control Point
cDNA	complementary deoxyribonucleic acid
CEN	European Committee for Standardization
CFU	colony forming units
DI	discrimination index
DIN	Deutsches Institut für Normung
DNA	deoxyribonucleic acid
EHEDG	European Hygienic Engineering and Design Group
ELGI	European Lubrication Grease Institute
EPDM	ethylene propylene diene monomer rubber
FDA	American Food and Drug Administration
GHP	Good Hygiene Practice
GMP	Good Manufacturing Practice
НАССР	Hazard Analysis Critical Control Point
HDPE	high density polyethylene
ISO	International Standardization Organisation
LMBA	Listeria monocytogenes Blood Agar

ME	microbicidal effect				
MPN	most probable number				
NBR	nitrile butyl rubber				
NLGI	National Lubricating Grease Institute				
OCP	own-checking plan				
PALCAM	Polymyxin Acriflavine Lithium Chloride Ceftazidime Aesculin				
	Mannitol agar				
PC	polycarbonate				
PCR	polymerase chain reaction				
PFGE	pulsed-field gel electrophoresis				
PMP	Pathogen Modeling Program				
PP	polypropylene				
PU	polyurethane				
PVC	polyvinyl chloride				
QAC	quaternary ammonium compound				
rRNA	ribosomal ribonucleic acid				
RT	ribotype				
RTE	ready-to-eat food				
SD	standard deviation				
TSB	Tryptone Soya Broth				
USDA	United States Department of Agriculture				
UVM	University of Vermont broth				
WTO	World Trade Organization				

1. Introduction

1.1 Equipment hygiene in food processing

Several factors have been reported to affect on the hygiene level of food processing equipment, including hygienic design of the equipment, hygienic practices of personnel, cleaning and disinfection of the equipment, lubricants used in the equipment as well as lay-out of the processing, air-currents, type of food product and cleanliness of the processing environment (Lelieveld et al., 2003). In this thesis, the first four factors were studied more specifically, with focus on occurrence of the pathogenic bacterium *Listeria monocytogenes* – a problematic contaminant in food processing.

1.1.1 Hygienic design of food processing equipment

Hygiene problems in equipment are caused when micro-organisms attach to the surfaces, survive on them and later become detached thus contaminating and reducing the quality of the product (Wirtanen, 1995). This can be due to a poor hygienic design in cases where the machines cannot be cleaned properly. Constructions that cause problems include sites where soil, product debris and micro-organisms can accumulate, e.g. dead ends, sharp corners and low-quality seals and joints (Anon., 1993; Anon., 1995). Equipment has been identified as the source of contamination in the food industry in many studies (see e.g. Table 4). In Finland hygiene in dairy plants was investigated by the Finnish Food Safety Authority in 2001–2005. Deficiencies were reported e.g. in topics concerning equipment hygiene as presented in Table 1. Additionally, L. monocytogenes was found in 313 (40-83 per year)/13876 (2342-4242 per year), i.e. 2.3% (1.7-2.9%) of the samples, which were taken from the processing equipment and the working environment (NFA, 1998, 2000-2005). Good hygienic design of food processing equipment protects the product from contamination with substances harmful to consumer health and provides access for cleaning, maintenance and inspection (Lelieveld et al., 2003). Criteria for good hygienic design are presented in Table 2.

Table 1. Deficiencies affecting on equipment hygiene observed in Finnish dairy plants during 2001–2005, inspected by the Finnish Food Safety Authority (397–450 inspections/year) (NFA, 1998, 2000–2005).

Торіс	No. of lack	s in inspections ^a (%)
Constructions and surfaces in the plant; type or wear of materials	Range: Total:	21–29 (4.7–6.7) 126 (5.8)
Layout of production rooms and arrangements of production	Range: Total:	12–17 (3.0–3.8) 74 (3.4)
Processing equipment and methods, temperature control	Range: Total:	1–10 (0.2–2.3) 30 (1.4)
Personnel: hair, clothing, jewellery, make-up, practices	Range: Total:	2–7 (0.5–1.6) 23 (1.1)
Cleaning and disinfection	Range: Total:	6–8 (1.4–1.8) 14 (1.6)

^a No. of plants inspected were 132 (2001), 132 (2002), 131 (2003), 126 (2004), 124 (2005).

1.1.2 Hygiene of personnel working with equipment

An important factor affecting the hygiene level of food processing equipment is the hygienic practices of people working with the equipment. Many papers have been published on unsatisfactory hygienic practices in food handling (Upmann and Reuter, 1998; Haupt et al., 1999; Henroid and Sneed, 2004) and outbreaks caused by these (Guzewich and Ross, 1999; Duncanson et al., 2003; LaPorte et al., 2003). Contamination of foods by food handlers has been identified as one of the most important causes of foodborne outbreaks (FDA, 2000; Michaels et al., 2004). One group of employees who frequently work with the equipment are the maintenance personnel, who dismantle machinery for cleaning procedures and reassemble it after cleaning in addition to maintaining the operation of machinery during production. No earlier reports are available on the effect of this specific group of personnel on production hygiene.

Personnel are both reservoirs and vectors of micro-organisms (Marriott, 1999; Holah and Taylor, 2003). The level and risk of contamination from personnel is difficult to measure as it depends on various factors such as the different activities and the range of personnel movement patterns during the working day as well as the perceptions and attitudes of the personnel (Troller, 1993; Holah and Taylor, 2003).

Table 2. Criteria for good hygienic design of food processing equipment (Anon., 1993; Anon., 1995; CEN, 1997; Lelieveld et al., 2003).

Design parameters	Generally recommended criteria for the food area ^a in equipment				
Construction materials	durable, cleanable, disinfectable; resistant to cracking, abrasion and corrosion; non-toxic, non-absorbent; do not transfer undesirable odours etc.; do not contribute to contamination of food. Suitable materials are e.g. stainless steels EN 1.4301 (AISI 304), EN 1.4404 (AISI 316L), EN 1.4435 (AISI 316I), EN 1.4571 (AISI 316Ti) and plastics (see Conveyer belts)				
Surface finish	cleanable, disinfectable, smooth, continuous, prevents trapping of microbes, $R_a \geq 0.8 \mu m$				
Drainability	self draining				
Corners	rounded, no dead spaces, cleanable, disinfectable				
Joints sealed, hygienic, no gaps or crevices, protruding ledges and seals sho avoided					
Welds	smooth, continuous; no misalignments, cracking or porosity; sloped edges				
Fasteners (screws, bolts)	avoid if possible; cleanable, disinfectable				
Seals/Gaskets	tolerate processing conditions without changes, cleanable, disinfectable, suitable materials include e.g. EPDM ^b , NBR ^c , nitrile rubber, silicone rubber, Viton rubber				
Rims	no ledges where product can lodge; cleanable, top rims rounded and sloped				
Bearings, shafts	located outside the food area, cleanable and disinfectable, food grade lubricant used				
Panels, covers, doors	prevent entry of soil and contaminants, cleanable, disinfectable				
Instrumentation and control devices	prevent ingress of contamination, sanitary couplings				
Conveyer belts	non-absorbent, covered edges, rounded rims, cleanable, disinfectable, tolerant; suitable materials PP^d , PVC^e , acetal copolymer, PC^f , $HDPE^g$				
Placing and installation	electronic devices in the non-food area, sealed to floor, rounded pedestal, clear space everywhere around the equipment to enable cleaning				

^a Area composed of surfaces in contact with food; the food area also includes the surfaces with which the product may come into contact under intended conditions of use, after which it returns into the product (CEN, 1997).

^b ethylene propylene diene monomer	^d polypropylene	^f polycarbonate
^c nitrile butyl rubber	^e polyvinyl chloride	^g high density polyethylene

Micro-organisms in the human body can be divided into resident and transient microbes. The face, neck, hands and hair contain both a higher proportion of transient micro-organisms and a higher general bacterial density than other parts of the body (Troller, 1993; Holah and Taylor, 2003). Hands are the major source of infection from transient and resident micro-organisms. *L. monocytogenes* has been found on hands or gloves used in food-handling (Destro et al., 1996; Autio et al., 1999). Hand hygiene has been considered to be the most important single operation affecting reduction of hygiene risk (Paulson, 2000; Holah and Taylor, 2003). It has also been estimated that approximately 5% of healthy people are asymptomatic carriers of *L. monocytogenes* in their intestines (Farber and Peterkin, 1991; Farber and Harwig, 1996; Skinner, 1996). Of the employees of poultry houses and slaughterhouses as many as 10–30% have been shown to be carriers (Bennett, 1986).

Contamination from personnel can be direct or indirect. Direct contamination can occur by contact between the body and the food product. In indirect contamination, people act as vectors and transfer contamination from one area or surface to another by e.g. using the same equipment in raw and cooked product areas (Holah and Taylor, 2003). Clothing or footwear can contaminate work surfaces when personnel move around the plant (Holah and Taylor, 2003). Several investigations on food-handling practices of personnel in hospitals (Angelillo et al., 2000; Askarian et al., 2004; Danchaivijitr et al., 2005) and food service environments (Henroid and Sneed, 2004; Sneed et al., 2004; Tang and Wong, 2004) have been published. Less information is available on hygienic practices of food employees in food plants or abattoirs, but several deficiencies in hygienic practices have been identified in the studies performed (Table 3).

Important means of avoiding contamination of products and equipment include Good Hygiene Practices (GHPs) and regular medical screening of the personnel, sufficient training of hygiene aspects and control of indirect contamination (Marriott, 1999; Holah and Taylor, 2003). The regulation on food hygiene (EU) No. 852/2004, Annex II (Anon., 2004a) state that food -handlers must be supervised and instructed in food hygiene matters commensurate with their work activity. Even if the personnel maintaining the equipment do not actually touch the raw materials or food products, they probably touch a multitude of surfaces in contact with unpacked products while maintaining the machinery.

	I HOLE J. DIMMES ON MY.	Sichie wormes practices and annuaces on	Joon upgicute of Joon manany personner.	
Food sector and country	Type of survey	Main conclu	tsions of hygienic practices	Reference
f mmos		Positive	Negative	
10 Meat plants, Austria	Survey of self-assessment		 relative frequency of non-conformities regarding own-checking was 25% at best, for premises/equipment/tools 10%, personal hygiene 21%, specific process hygiene 30% and for cutting plant hygiene 9% at best. Simple hygiene fundamentals were not used. 	Haupt et al., 1999
54 Abattoirs, the Netherlands	Inspection with checklist, microbiological sampling, two visits	+ during the follow-up study improved consciousness of Good Manufacturing Practices (GMPs) among employees was observed	- better consciousness of importance of GMPs required, e.g. more frequent hand washing	Heuvelink et al., 2001
5 Abattoirs, Finland	Visual observation and microbiological sampling	+ frequency of slaughtering errors and touching of the carcasses was low	 designing slaughtering lines so as to make hygienic working possible and the enforcement of hygienic practices, such as the regular disinfection of working tools, were needed. Hygienic practice was associated with the carcass contamination level. 	Rahkio and Korkeala, 1996
Meat abattoir, Austria	Visual observation		 defects in personnel clothing and behaviour, personal belongings stored in slaughter room 	Upmann and Reuter, 1998
Abattoir, South Africa	A structured questionnaire	+ basic hygiene practices were used and the employees adhered to majority of them	 improved communication between management and workers and more training needed 	Nel et al., 2004
Small food manufacturing plants, UK	Structured interviews with managers on hygiene training	+ all the firms had taken some hygiene training, managers had positive attitudes towards training	 hygiene programmes unstructured and unrecorded, regular refresher training given in less than half of the plants, lack of time and money were perceived to be a major restricting factors to training. 	Worsfold, 2005
Bakery, USA	Microbiological sampling		 food handler was a source of viral contamination in baked goods 	Weltman et al., 1996
Flamenquin plant, Spain	Microbiological sampling		 raw materials and food handlers were the principal sources of microbiological contamination 	Cordoba et al., 1998
Food industry (meat, dairy, bakery, fish), Finland	Mail survey on attitudes towards own-checking and HACCP plans was distributed to 87 plants, with response rate of 34.9%	 + attitudes exclusively positive, all 30 companies had a functioning own-checking plan (OCP), while other quality management programs were less prevalent 	 most difficulties in devising the OCP/HACCP plan caused choosing the critical control points, committing the firm's entire workforce and organizing the documentation 	Hielm et al., 2006
Food handlers in general, Italy	Interview with a structured questionnaire within a random sample from 411 food handlers	+ positive attitude toward foodborne disease control by the great majority, more likely achieved when more training received	 the attitude was not supported by hygienic practices, e.g. only 20.8% used gloves when touching unwrapped raw food, strong need for educational programs for improving knowledge 	Angelillo et al., 2000
20 Food premises, UK	An audit proforma, questionnaires on effectiveness of food hygiene training		 formally trained staff work more hygienically than non-trained 	Kirby and Gardiner, 1997
52 Small and medium size (SME) food businesses, UK	Questionnaires on food handler (including food handlers working at retail, catering or, manufacturing) beliefs and self-reported practices on food safety	+ food handlers were aware of the food safety actions they should be carrying out, 95% had received food hygiene training (100% in the manufacturing sector). Food handlers in the manufacturing sector were more likely to report carrying out food safety behaviours than were more likely to report carrying out food safety behaviours than were more likely in the catering and retail sector.	 - 63% admitted to sometimes not carrying out food safety procedures, lack of time, personnel, resources 	Clayton et al., 2002

Table 3. Studies on hysienic working practices and attitudes on food hysiene of food industry personnel.

1.1.3 Lubricants used in equipment

The purpose of the use of lubricants in food-processing equipment is to reduce friction and wear, inhibit the access of outside particles to equipment surfaces, protect surfaces against corrosion and remove wear particles, increase the efficiency of systems and the transfer of heat, power and electricity (Netuschil, 1995; Lewan, 2003). The lubricants usually used in the food industry are composed of base oil (e.g. mineral oil, white oil, silicone), thickeners and additives. The lubricants can be completely oil-based but many are emulsions containing water. To achieve high technical performance, the lubricants are often made of different synthetic components (Netuschil, 1995).

Lubricants can be contaminated with water, organic material, residues of other lubricants, physical or chemical substances causing oxidation and other chemical reactions, particles from corrosion (Anon., 2003), or with micro-organisms. Contamination in lubricants can lead to contamination of food products e.g. through leakage from bearings, dripping from open lubrication points e.g. chains, leakage from oil circulation systems or from corroded joints of oil-filled heat exchange systems or contact between oil-coated machine surfaces (Anon., 2003). The microbes must often tolerate anaerobic conditions and low water activity in lubricants. L. monocytogenes is capable of withstanding the above mentioned conditions (Buchanan et al., 1989; Lou and Yousef, 1999). It has been shown to survive in butter, which was the vehicle in a Finnish L. monocytogenes epidemic in 1998–1999 (Lyytikäinen et al., 2000). Rossmoore (1988) reported findings of *L. monocytogenes* in dairy conveyer lubricants. Use of lubricants in conveyers has also caused hygiene problems in breweries (Heinzel, 1988). Acinetobacter sp., Algaligenes sp., Pseudomonas sp. and sulphate reducing bacteria have been isolated from lubricants (Ortiz et al., 1990; Hamilton, 1991). Petitdemange et al. (1995) observed clear differences between strains of *Clostridium butyricum* in their ability to survive and grow in industrial glycerol.

During processing and cleaning the quality of lubricants is impaired. The survival of microbes in lubricants has been reported to be enhanced when the lubricants are contaminated with organic material and water (Netuschil, 1995; Lou and Yousef, 1999). The addition of antimicrobial substances such as glutaraldehyde (25 ppm) or isothiazoline (10 ppm) has been reported to inhibit the growth of microbes, e.g. *L. monocytogenes* (Rossmoore, 1988; Hsu, 1991).

1.1.4 Listeria monocytogenes in equipment hygiene

L. monocytogenes is a bacterium of special concern in food processing as it has been connected to several food-borne outbreaks and it causes listeriosis, a severe illness especially for immunocompromized individuals, with high mortality of approximately 20–30% (Rocourt, 1996; Rocourt et al., 2003; Lyytikäinen et al., 2006). The bacterium causes problems in food processes, as it tolerates well various stress factors that it encounters in food processing plants and also has a very good ability to attach to different surfaces and thus to persist in the food plants for years. *L. monocytogenes* has been detected from food processing equipment which has been implicated in the contamination of final products (Table 4). It has been isolated from the equipment and processing environments of various food sectors in several studies (Gravani, 1999; Tompkin, 2002). The characteristics of this bacterium are presented in detail in Section 1.3.

1.2 Legislation and standards

1.2.1 Legislation on hygienic design of food processing equipment

In Europe the most important legislation giving criteria for hygienic design of equipment is the Council Directive on the approximation of the laws of Member States relating to machinery (89/392/CEE, revised 98/37/EC, Anon., 1998). It contains safety requirements and a few very basic principles of hygienic design on constructions and surfaces, which must be cleanable and safe for production. This directive also obliges the equipment manufacturers to provide guidelines on sanitizing the equipment. EC Regulation No. 1935/2004 on materials and articles intended to come into contact with food applies to materials and articles which already are or are intended to come into contact with food (Anon., 2004b). According to this regulation the materials shall not transfer their constituents to food in quantities which could endanger human health or cause changes in composition or organoleptic characteristics of food. There is also a EC Regulation No. 2023/2006 on good manufacturing practice for materials and articles intended to come into contact with food (Anon., 2006).

Equipment	Food sector and	product		С	ountry	7	R	eference		
been associated with contamination of final product.										

Table 4. L. monocytogenes findings in food processing equipment which have

Equipment	Food sector and product	Country	Reference
Air chiller	Poultry, raw	Finland	Miettinen et al., 2001
Cold-smoker	Fish, cold-smoked salmon	USA	Norton et al., 2001
Conveyer belts	Meat	Nordic country	Suihko et al., 2002
	Seafood	Nordic country	Suihko et al., 2002
	Poultry, raw	Finland	Miettinen et al., 2001
Dicing machine	Meat, cooked	Finland	Lundén et al., 2002
Freezer	Poultry, cooked	USA	Berrang et al., 2002
Packaging machine	Dairy, butter	Finland	Maijala et al., 2001
	Dairy, ice cream	Finland	Miettinen et al., 1999a
Salting machine	Fish, cold-smoked salmon	Finland	Autio et al., 1999
	Fish, cold-salted, cold-smoked and smoked salmon	Finland	Johansson et al., 1999
Skinning machine	Meat	Nordic country	Suihko et al., 2002
	Poultry, raw	Finland	Miettinen et al., 2001
	Fish, cold-salted, cold-smoked and smoked salmon	Finland	Johansson et al., 1999
Slicing machine	Fish, cold-smoked salmon	Finland	Autio et al., 1999
	Fish, cold-salted, cold-smoked and smoked salmon	Finland	Johansson et al., 1999
	Meat	Nordic country	Suihko et al., 2002
- slicer switches	Meat, rillets	France	Goulet et al., 1998
	Meat	_ ^a	Lelieveld et al., 2003
Tumbling machine	Meat, cooked	Greece	Samelis and Metaxopoulos, 1999
Evisceration machine, pluck sorter, spin chiller	Poultry, raw	Denmark	Ojeniyi et al., 1996

^a not reported

A basic standard concerning hygiene requirements for the design of machinery is the International Standardization Organisation (ISO) standard 14159:2002 (ISO, 2002). It specifies hygiene requirements of machines and user information to be provided by the manufacturer. It applies to all types of machines and associated equipment used in applications where hygiene risks to the consumer of the product can occur. The European Committee for Standardization (CEN) issues standards for equipment manufacturers to enable fulfilling the requirements of the EU directive. One important basic standard is the standard EN 1672-2 "Food processing machinery - Safety and hygiene requirements - Basic concepts - Part 2; Hygiene requirements" (CEN, 1997). There are also a number of standards for different specific food processing equipment, e.g. for slicing, cutting and filling machines (http://www.cen.eu/; http://www.3-a.org). Guidelines and methods in

accordance with the standards are published by different organizations, e.g. by the European Hygienic Engineering and Design Group (EHEDG) (<u>http://www.ehedg.org</u>), 3-A Sanitary Standards Inc. (<u>3-A, http://www.3-a.org</u>) and NSF International, The Public Health and Safety CompanyTM (<u>http://www.nsf.org</u>).

1.2.2 Legislation on lubricants used in food processing

Lubricants used in food processing equipment at points where incidental contact with food products may occur (i.e. food-grade lubricants), have special requirements. They must fulfil the requirements of legislation and be internationally accepted, physiologically safe and neutral in taste and odour (Netuschil, 1995; Köhler, 2001). Use of food-grade lubricants is recommended in food-processing plants especially at critical control points (Anon., 2003). Definitions of food-grade lubricants can be found in the document FGL1/2001/issue 2 of the European Lubrication Grease Institute (ELGI), the National Lubricating Grease Institute (NLGI) and the EHEDG. The standard of Deutsches Institut für Normung (DIN) V 0010517 and the NSF International draft for an American National Standards Institute (ANSI) standard are also sources for official information on lubricants (Anon., 2003). These definitions include the registration of food-grade lubricants (H-1) by a competent third party organisation, as also required in the new ISO standard "Safety of machinery -Lubricants with incidental product contact – Hygiene requirements" (ISO, 2006). This standard specifies definitions and hygiene requirements for the formulation, manufacture, use and handling of lubricants which may come into incidental contact with products during manufacture and processing. The United States Department of Agriculture (USDA) has classified the lubricants into foodgrade and non-food-grade. USDA H1 lubricants contain only components approved by The American Food and Drug Administration (FDA) and they can be used in places in which there is incidental contact with food. These classifications are also generally accepted in Europe, where legal requirements are less stringent (Köhler, 2001). The USDA stopped registering lubricants in 1998 and these records are nowadays maintained by NSF International (Yano, 2005). Additionally, a guideline on "Production and use of food-grade lubricants" has been published by EHEDG (Anon., 2003).

1.2.3 Legislation on L. monocytogenes

According to EC Regulation No. 2073/2005 on Microbiological criteria for foodstuffs (Anon., 2005), the limit of L. monocytogenes is 100 CFU/g at the end of the shelf-life of ready-to-eat (RTE) food products. In the same regulation it is also stated that food businesses manufacturing RTE foods, which may pose a L. monocytogenes risk for public health, shall sample the processing areas and equipment for L. monocytogenes as part of their sampling scheme (Anon., 2005). The presence of *L. monocytogenes* in meat and fish products is not regulated by Finnish food legislation, but in recent years several guidelines on control of Listeria in the food chain, targeted at meat and fish processing facilities and retailers, have been published by the authorities (http://www.evira.fi, http://www.ktl.fi). The National Food Agency of Finland has given press releases on vacuum-packed fish products, advising consumers to pay attention to the time for which the products are stored; to the home refrigerator temperature, which should be below 3 °C; and to consuming products before the best-by-date (Lyytikäinen et al., 2006). In the United States there is currently a zero tolerance policy for the levels of L. monocytogenes allowed in food (McLauchlin et al., 2004). In Canada, the limit is 100 CFU/g in RTE foods (Health Canada, 2004).

1.3 Characteristics of *L. monocytogenes* – a problematic food processing contaminant

1.3.1 The genus Listeria and L. monocytogenes

The genus *Listeria* consists of six species and two subspecies: *L. monocytogenes*, *L. ivanovii* subs. *ivanovii*, *L. ivanovii* subsp. *londoniensis*, *L. seeligeri*, *L. innocua*, *L. welshimeri* and *L. grayi*. Two of the species are pathogenic: *L. monocytogenes*, the foodborne human pathogen and *L. ivanovii*, an animal pathogen. *L. ivanovii* and *L. seeligeri* have also occasionally been associated with human listeriosis (Cummings et al., 1994; Lessing et al., 1994; Khelef et al., 2006).

Bacteria belonging to the genus *Listeria* are Gram-positive and non-sporeforming. They are catalase-positive, oxidase-negative and methyl red-positive bacteria. Voges-Proskauer reaction is positive. They are able to move with the aid of flagella at 20–25 °C. The bacteria are regular, short rods; some of the cells may be curved. Their diameter is 0.4–0.5 μm and length 0.5–2 μm (Seeliger and Jones, 1986).

L. monocytogenes was first described by Murray et al. (1926) and was then named as *Bacterium monocytogenes* because it caused severe monocytosis in infected laboratory rabbits and guinea pigs. Later, in 1927 Pirie renamed the bacterium as *Listerella hepatolytica* and further in 1940 to its current name (Gray and Killinger, 1966).

1.3.2 Occurrence

Listeria, including *L. monocytogenes*, are ubiquitous in nature and can be found in water, mud, sewage and vegetation as well as in cattle milk, faeces of animals and humans, animal feed as well as in food and almost all food raw materials (Seeliger and Jones, 1986; Bell and Kyriakides, 2004; Khelef et al., 2006).

L. monocytogenes has been isolated from a large variety of both raw and cooked food products including meat products e.g. raw beef, lamb, pork, ground meats, ham, fermented and dried sausages, processed meats and pâté; poultry products e.g. raw and cooked poultry and eggs; sea food products such as raw fish, smoked salmon, uncooked and cooked shellfish; dairy foods e.g. raw and pasteurized milk, creams, soft, semisoft and hard cheeses and ice cream; vegetables including cabbage, cucumber, potatoes, tomatoes and frozen vegetables; and RTE foods (Farber and Peterkin, 1991; Bell and Kyriakides, 2004). The bacterium was recognized as a foodborne pathogen after several outbreaks in the early 1980s (Schlech et al., 1983; Bell and Kyriakides, 2004). Food products which have caused outbreaks of listeriosis include a variety from different areas of the food industry: cole slaw in Canada 1981 (Schlech et al., 1983), pasteurized milk in USA 1983 (Fleming et al., 1985), Mexican style cheese in USA 1985 (Linnan et al., 1988), soft cheese in Switzerland 1983–1987 (Bille, 1990), pâté in UK 1987–1989 (McLauchlin et al., 1991), jellied pork tongue in France 1992–1993 (Goulet et al., 1993), chocolate milk in USA 1994 (Dalton et al., 1997), corn and tuna salad in Italy 1997 (Aureli et al., 2000), rainbow trout in Sweden 1994–1995 (Ericsson et al., 1997) and in Finland 1997 (Miettinen et al., 1999b), delicatessen turkey meat in USA 2000 (Olsen et al., 2005), 2001 (Frye et al., 2002) and 2002 (Gottlieb et al., 2006), as well as butter in Finland 1998–1999 (Lyytikäinen et al., 2000). Cases have also been caused by e.g. uncooked chicken meat (Schwartz et al., 1988; Anon., 1989; Kaczmarski and Jones, 1989). Summaries on the occurrence of this pathogen worldwide in different food products have been published (Farber and Peterkin, 1991; Autio, 2003) and therefore in this work a summary of the occurrence and amounts of *L. monocytogenes* in various food products in Finland, where the studies of the current thesis were also performed, is presented in Table 5.

1.3.3 Tolerance of pH, temperature and a_w

L. monocytogenes is able to survive in various environmental conditions. It is a facultatively anaerobic bacterium, which grows well in aerobic conditions (Khelef et al., 2006). It grows over a wide pH-range of 4.3–9.6 (Lou and Yousef, 1999), optimal growth being at neutral to slightly alkaline pH (Seeliger and Jones, 1986). Survival in as low pH as 1.4 has been reported (Reimer et al., 1988).

The temperature growth range of *L. monocytogenes* is 1–45 °C (Seeliger and Jones, 1986), but slow growth at as low as -0.4 °C (Walker et al., 1990) and even at -1.5 °C (Hudson et al., 1994) have been reported as well as good survival in much lower freezing temperatures (Lou and Yousef, 1999), e.g. many weeks in foods at -18 °C (Golden et al., 1988; Olsen et al., 1988). The optimum growth temperature is 30–37 °C (Seeliger and Jones, 1986). As the bacterium grows in refrigeration temperatures, it is a potential risk in cold rooms of food processing plants (Gravani, 1999). Cold stress adaptation of *L. monocytogenes* is a function of many molecular adaptation mechanisms and is an important characteristic of *L. monocytogenes*, enabling it to survive and proliferate in refrigerated foods and cold environments (Tasara and Stephan, 2006).

Table 5. Occurrence and amount of L. monocytogenes (L.m.) *in various food products in Finland during 1996–2005.*

Food product	No. of <i>L.m.</i> positive samples / analyzed samples (%)	Quantity of <i>L.m.</i> (CFU/g): no. of samples / no. of all samples (%)	No. of producers with <i>L.m.</i> (%)	Year of study	Reference
	<u>Fi</u>	sh and other seafood			
cold-salted fish; - vacuum-packed	44/172 (25.6) 22/110 (20.0)	$\leq 100:10/16 (62.5)$ $\geq 100: 6/16 (37.5)$	16/46 (34.8)	1996–2000 1996	Johansson et al., 2003 Johansson et al., 1999
 rainbow trout, non-sliced rainbow trout, sliced 	4/12 (33.3) 10/31 (32.3)				Lyhs et al., 1998 Lyhs et al., 1998
- not vacuum-packed	28/204 (13.7) 0/24 (0) 13/81 (16.0)			2003-04	Aalto et al., 2006 Johansson et al., 2003 Aalto et al. 2006
cold-smoked fish; - vacuum-packed	44/223 (19.7) 46/356 (13)	<100: 32/43 (74.4) ≥ 100: 11/43 (25.6)	7/37 (18.9)	2003-04 2003-04 2001	Aalto et al., 2006 Hatakka and Johansson, 2002
	76/720 (10.6) 5/30 (17)	$\leq 100: 1/5 (20.0)$ >100: 4/5 (80.0)	3/12 (25)	1996–2000 1996	Johansson et al., 2003 Johansson et al., 1999
- salmon - rainbow trout, non-sliced - rainbow trout, sliced	22/22 (100) 4/42 (9.5) 5/20 (25.0)		1/1 (100)	1997	Autio et al., 1999 Lyhs et al., 1998 Lyhs et al., 1998
cold-smoked fish; not vacuum- packed	4/54 (7.4)			2003–04	Aalto et al., 2006
cold-salted and cold-smoked fish	1/10 (10.0)			1996-2000	Johansson et al., 2003
- vacuum-packed - not vacuum-packed hot smoked fish; - vacuum-packed - rainbow trout	6/369 (16.3) 6/42 (14.3) 2/147 (1.4) 1/48 (2) 1/42 (2.4)	≤ 100: 1/1 (100.0)	1/12 (8.3)	1996–1999 1996–1999 1996–2000 1996	Anon., 2000b Anon., 2000b Johansson et al., 2003 Johansson et al., 1999 Lyhs et al., 1998
- salmon hot smoked fish - not vacuum-packed roe	0/6 (0) 0/8 (0) 0/29 (0) 7/147 (4.8) 2/18 (17)		0/11 (0) 7/26 (26.9)	1996–2000 2003–04 1999	Lyhs et al., 1998 Johansson et al., 2003 Aalto et al., 2006 Miettinen et al., 2003 Johansson et al., 2003
crustaceans and monuses	5/18(17)	Dairy		1990–2000	Johansson et al., 2005
soft and semisoft cheeses from					
pasteurized milk soft and semi-soft cheeses from raw	0/132 (0)			2003–04	Aalto et al., 2006
milk soft cheese unripened cheese	5/90 (5.5) 0/49 (0) 2/144 (1.4)	$1.1 - 6 \times 10^3$		2003–04 1996–2000 1996–2000	Aalto et al., 2006 Johansson et al., 2003 Johansson et al., 2003
milk and milk products	6/842 (0.7)			2001	Hatakka and Maijala, 2003
ice cream ice creams and desserts	6/1129 (0.5) 0/193 (0)		1/1 (100)	1990–1997 2001	Miettinen et al., 1999a Hatakka and Maijala, 2003
milk products	43 (1–19/year)/ 15476 (1531– 2961/year), i.e.			1998, 2000–2005	(NFA, 1998, 2000–2005)
butter	34/59 (57.6)	$\leq 100: 16/18 (88.9)$ 2/18 (11.1)	1/1 (100)	1998–1999	Maijala et al., 2001
edible fats and oils		(****)	0/27 (0)	2001	Hatakka and Maijala, 2003

Table 5. Cont.

Food product	No. of <i>L.m.</i> positive samples / analyzed samples (%)	Quantity of <i>L.m.</i> (CFU/g): no. of samples / no. of all samples (%)	No. of producers with <i>L.m.</i> (%)	Year of study	Reference
Meat and meat products					
meat and meat products	128/857 (14.9)			2001	Hatakka and Maijala, 2003
cold cuts	0/236 (0) 0/30 (0)		0/4 (0)	2002 1996–2000	Johansson et al., 2002 Johansson et al. 2003
pieces of raw broiler	35/100 (35)			1996-2000	Johansson et al., 2003
	38/61 (62.0)	5.01×10^{3}	3/3 (100)	1997–1998	Miettinen et al., 2001
	4//85 (55.5)	MPN/kg ^a	4/4 (100)		Husu et al., 1995
– frozen carcasses	27/80 (33.8)	≤ 2250 CFU/carcass			Johansson and Hirn, 1992
eggs and egg products	0/6 (0)			2001	Hatakka and Maijala, 2003
Vegetables and fruits					
fresh vegetables and products	2/158 (1.3)			1996-2000	Johansson et al., 2003
frozen vegetables	61/313 (19.5)			1996-2000	Johansson et al., 2003
fruits and vegetables and products	9/452 (2.0)			2001	Hatakka and Maijala, 2003
berries	4/344 (1.2)			1996–2000	Johansson et al., 2003
Others					
RTE foods	2/78 (2.6)			1996-2000	Johansson et al., 2003
	6/119 (5.0)			2001	Hatakka and Maijala, 2003
soups and sauces	0/88 (0)			2002	Hatakka and Maijala, 2003
confectioneries	1/23 (4.3)			1996-2000	Johansson et al., 2003
salted mushrooms	1/1(100)	3.8×10^{6}			Junttila and Brander, 1989
grain and grain products	0/11 (0)			2001	Hatakka and Maijala, 2003
nuts, nut products and snacks	0/2 (0)			2001	Hatakka and Maijala, 2003
herbs and spices	0/1 (0)			2001	Hatakka and Maijala, 2003

^a MPN = most probable number

L. monocytogenes does not survive heating at 60 °C, 30 min and thus is destroyed in pasteurization treatment (Seeliger and Jones, 1986). However, if the heating process does not reach all the material to be treated, or if the amount of bacterial cells is very high, e.g. 10^5 – 10^6 CFU/ml, the bacterium may survive (ICMSF, 1996). Tolerance against heating may also vary depending on several factors (grease content, NaCl-content, atmosphere, heating profile, strain) (Embarek and Huss, 1993). Unusual tolerance of *L. monocytogenes* to thermal processing in chicken/broiler meat has been observed in some studies, e.g. as high as 82.2 °C internal temperature was not sufficient to kill *L. monocytogenes* when inoculated on raw chicken meat surface at a level of 10^5 – 10^6 CFU/g

bacteria (Carpenter and Harrison, 1989). Thorough and sufficient heat treatment (e.g. 85 °C, 15 min or 80 °C, 20 min) is generally an effective way to destroy *L. monocytogenes* from cleaned production surfaces (Marriott, 1999). However, grease has been suggested to protect *L. monocytogenes* from heat, at least in some food products (Embarek and Huss, 1993; Murphy et al., 2004).

L. monocytogenes is able to grow at water activity (a_w) values of ≥ 0.90 (Farber et al., 1992; Lou and Yousef, 1999). It tolerates well high salt concentrations (>20%) and thus it can cause problems in brine solutions and brining machines in food processing (Gravani, 1999). The nutritional requirements of *L. monocytogenes* are similar to those of many other gram-positive bacteria (ICMSF, 1996; Jay, 1996).

1.3.4 Listeriosis

Listeriosis is an illness which is severe especially to individuals belonging to risk groups, i.e. very young or old people, pregnant women, cancer and AIDS patients and patients receiving immunosuppressive therapy (Rocourt, 1996). Approximately 20% of the population typically belong to risk groups (Miller et al., 1997; Maijala et al., 2001). The fatality rate of the disease is approximately 20–30% (Rocourt, 1996; Rocourt et al., 2003; Lyytikäinen et al., 2006). The main symptoms include miscarriages in pregnant women and meningitis and bacteraemia in neonates and adults (Rocourt, 1996). However, milder food poisoning symptoms including vomiting and diarrhoea are possible, or in case of zoonotic infection localised skin lesions. The disease can also be asymptomatic or very mild, but may later develop into infections such as meningitis. The incubation time of the disease varies from <24 h to several months (Bell and Kyriakides, 2004). The factors predisposing infection are not fully understood, but include host immunity, level of inoculum and virulence of the strain (Bell and Kyriakides, 2004).

L. monocytogenes has 13 serovars based on the expression of somatic (O) and flagellar (H) antigens. Approximately 95% of human isolates belong to serovars 4b, 1/2a and 1/2b (Graves et al., 1999). Invasive outbreaks have mainly been caused by serovar 4b (Slutsker and Schuchat, 1999). *L. monocytogenes* is transmitted to humans via three main routes; contact with animals, cross-infection

of new-born babies in hospital, and foodborne infection, of which the last two routes apply to the majority of human listeriosis cases (Bell and Kyriakides, 2004).

In Finland 18–53 cases of listeriosis have been reported annually during 1995–2004 (Lyytikäinen et al., 2006) and have been connected e.g. to consumption of salted mushrooms (Junttila and Brander, 1989), rainbow trout (Miettinen et al., 1999b) and butter (Lyytikäinen et al., 2000). Almost 25% (78/315) of listeriosis cases in Finland have been caused by a certain sero-genotype or closely related genotypes, which have also been found from vacuum-packed cold-smoked or cold-salted (i.e. 'gravad') fish products, which accordingly are considered risk products especially to people belonging to risk groups (Lyytikäinen et al., 2006).

For healthy adults, the doses of *L. monocytogenes* causing listeriosis have been reported to vary from 10^5 to 10^9 CFU/g or /ml (Junttila and Brander, 1989; Misrachi et al., 1991; Dalton et al., 1997; Miettinen et al., 1999a, b; Aureli et al., 2000). However, for risk groups the doses have been reported to be lower, varying from <10 to 10^4 CFU/g (Berrang et al., 1988: Ericsson et al., 1997; Anon., 2000a). Despite the recent emphasis on studying this bacterium, the sources and routes of contamination, as well as the infective dose have still in many cases remained unknown (Rocourt et al., 2003). Currently the opinion is that high levels of *L. monocytogenes* are needed for causing the illness in the normal population (Chen et al., 2003; FAO/WHO, 2004).

1.3.5 Detection, identification and typing

Direct plating, selective enrichment, cold enrichment and several rapid methods based on e.g. polymerase chain reaction (PCR) can be used in various combinations to detect *L. monocytogenes* in food, clinical and environmental samples. Several methods for typing of *L. monocytogenes* isolates are also available (Graves et al., 1999).

A typical method for isolation of *L. monocytogenes*, especially from food products, includes one- or two-step enrichment followed by plating on selective agar. Direct plating is also used in detection from contaminated or infected material (Khelef et al., 2006). Standards of different organizations are available and one of the most commonly used are the methods of the International

Standardization Organization (ISO), ISO 11290-1 for detection (ISO, 1996) and ISO 11290-2 (ISO, 1998) for enumeration. Typical enrichment broths used in enrichment of *L. monocytogenes* are e.g. Fraser-broth (Fraser and Sperber, 1988) and University of Vermont broth (UVM) (Donnelly and Baigent, 1986; ICMSF, 1996). The bacterium grows well in many common media, e.g. brain heart infusion and trypticase or tryptose broth (Jay, 1996). Suspected samples are cultivated on selective agars e.g. Oxford- (Curtis et al., 1989), Modified Oxford-(MOX) (Lee and McClain, 1986), or Polymyxin Acriflavine Lithium Chloride Ceftazidime Aesculin Mannitol -agar (PALCAM) after an enrichment step, (van Netten et al., 1989). Special agars have also been developed for identification of *L. monocytogenes*, such as a *Listeria monocytogenes* Blood Agar (LMBA) based on the β -hemolytic reaction on sheep blood and *Listeria* selective agents (Johansson, 1998).

Confirmation tests of *L. monocytogenes* are carried out after incubation on selective agar plates and strain purification on non-selective agar plates. The genus confirmation is typically based on gram-staining, determination of motility at 20–25 °C, biochemical tests e.g. catalase test and especially for *L. monocytogenes* also the β -hemolysis test. *L. monocytogenes* produces β -hemolysis on blood agar plates, with a narrow zone of hemolysis around colonies. However, *L. ivanovii* and *L. seeligeri* also produce hemolysis. Additional tests are based on utilization of different sugars (ICMSF, 1996). These tests can be carried out with commercial kits, e.g. API Listeria[®] test (Bio-Mérieux SA, Marcy l'Etoile, France).

In identifying *L. monocytogenes* -genotypes more than ten different molecular typing methods have been applied (Graves et al., 1999). Hitherto, the only fully automated typing method available is ribotyping (Bruce, 1996). Ribotyping refers to the use of nucleic acid probes to recognize ribosomal genes (Farber, 1996). In ribotyping DNA is first extracted from cells, digested with an endonuclease (restriction enzyme), e.g. *Eco*RI, followed by separation of fragments by agarose gel electrophoresis. Separated fragments are transferred to a nylon membrane and hybridized with a labelled cDNA-probe derived from rRNA by reverse transcriptase. A chemiluminescent pattern is created and recorded (Jay, 1996). In automated ribotyping, the system performs all the process steps required to characterize the isolated bacterial colony to the strain level, from cell lysis to image analysis (Bruce, 1996). The process contains the following steps, of which only the first and last are performed manually: sample

preparation from bacterial cultures, DNA preparation, DNA separation and transfer, membrane processing using a labelled 16s rRNA probe from E. coli, detection of the chemiluminescent fragments using a charge-coupled device (CCD) camera, analysis by comparison of the patterns and optionally data manipulation and sorting (Farber, 1996). The results for 32 strains processed in batches of 8 can be obtained in 24 h. Each strain produces a unique DNA fragment pattern, which the system uses in a series of proprietary algorithms to generate a RiboPrint pattern. The pattern is characterized, archived and compared to a supplied database as well as against all the other patterns which have been run on the system to determine similarity (Bruce, 1996). Manual (Baloga and Harlander, 1991; Graves et al., 1994; Jensen et al., 1996; Louie et al, 1996; Ojeniyi et al., 1996; Kerouanton et al., 1998) and automated (Ryser et al., 1996; Arimi et al., 1997; Wiedmann et al., 1997; Allerberger and Frischel, 1999; Gendel and Ulaszek, 2000; Norton et al., 2001; Suihko et al., 2002; Lukinmaa et al., 2004; Grif et al., 2006; Saunders et al., 2006) ribotyping have been used for typing of L. monocytogenes in the studies cited.

In PFGE the genomic DNA is digested by one or more restriction enzymes, the fragments are separated by field inversion electrophoresis and fragments are resolved in agarose gels. The alternating electrical fields force molecules to change directions, resulting in electrophoretic profiles, i.e. designated pulsovars (Jay, 1996). Many factors, such as electric field strength, field angle and shape, agarose type and concentration, pulse time, ionic strength and temperature, are known to affect the resolution of this highly discriminating and reproducible method (Farber, 1996). PFGE has proved to be a very accurate and reproducible method for molecular typing of *L. monocytogenes* (Brosch et al., 1996) and has been used in numerous studies on contaminants and in epidemiology (Destro et al., 1996; Autio et al., 1999; Graves et al., 1999; Lukinmaa et al., 2004; Miettinen and Wirtanen, 2006). PFGE is the standard method used in PulseNet, the molecular subtyping network for foodborne disease surveillance. Via PulseNet public health officials can share molecular epidemiological information in real-time (Swaminathan et al., 2006).

Serotyping is the traditional and still routinely used phenotyping method in cases of outbreaks. However, it has a relatively poor discrimination power and some industrial isolates may be untypeable with standard typing antisera (Graves et al., 1999). It is based on differing antigenic determinants expressed on the cell
surface. Antigenic variations are produced by many different surface structures, which can be identified by serological typing. In the case of *Listeria*, the strains are divided into serotypes based on somatic (O) and flagellar (H) antigens. *L. monocytogenes* has 13 different serovars (Seeliger and Jones, 1986).

1.3.6 Attachment and transfer

L. monocytogenes attaches to and grows on different kinds of surfaces even at low temperatures (Mafu et al., 1990; Wirtanen and Mattila-Sandholm, 1993; Smoot and Pierson, 1998) and forms biofilms (Jeong and Frank, 1994; Blackman and Frank, 1996). According to Mafu et al. (1990) *L. monocytogenes* attaches to stainless steel, glass and plastic surfaces both at 20 °C and 4 °C in 20 min or 1 h, respectively. Several factors affect attachment and biofilm formation of *L. monocytogenes*, such as the type of surface and the level of nutrients available (Ronner and Wong, 1993), the bacterial strain (Lundén et al., 2000; Norwood and Gilmour, 1999) as well as the presence of other micro-organisms (Sasahara and Zottola, 1993; Jeong and Frank, 1994). According to a study of Mai et al. (2006), the overall wettability of the surfaces appeared to be a primary determinant of attachment of *L. monocytogenes*. Also corrosion enhances the attachment (Mai et al., 2006).

Recontamination, i.e. transfer of pathogens to the product from the processing environment after an inactivation step (den Aantrekker et al., 2003), has been identified as a significant cause of contamination of foods and thereby foodborne illnesses (Reij and den Aantrekker, 2004). Recontamination is effected by transfer of contaminants from surface to surface. Transfer of persistent *L. monocytogenes* contamination between food-processing plants was associated with a dicing machine in the study of Lundén et al. (2002). Only a few studies have been made of the transfer of *L. monocytogenes* from production surfaces to products or vice versa. Studies reported, and predictive models available, are summarized in Table 6. These studies show that *L. monocytogenes* is readily transferred to different products from various processing surfaces. *Table 6. Studies on transfer of* L. monocytogenes *to food products and predictive models constructed.*

Transfer/cross-c	ontamination	Model	Reference
from	to	(yes/no)	
Stainless steel slicer blade	Turkey breast, bologna and salami; slicer surfaces	No	Vorst et al., 2006a
Stainless steel slicer blade	Uncured, oven roasted turkey, bologna, salami; slicer housing, conveyer belt	No	Lin et al., 2006
Stainless steel kitchen knifes	Turkey breast, bologna and salami; slicer surfaces	No	Vorst et al., 2006b
Stainless steel, PVC, polyurethane (PU)	Beef	Yes	Midelet and Carpentier, 2002
Stainless steel, high density polyethylene (HDPE)	Bologna and American cheese	No	Rodríguez and McLands-borough, 2007
Stainless steel, PVC and PU (pure and two-species biofilms)	Tryptone soya agar (TSA)	Yes	Midelet et al., 2006
Food contact surfaces, gloves, environment	Fish products	Yes	Ivanek et al., 2004
Food processing environment	Different foods	Yes	Schaffner, 2004
From inoculated oranges to work surfaces	From work surfaces to orange juice	No	Martinez-Gonzales et al., 2003
From contaminated utensils to uninoculated oranges	To orange juice		
Bare or gloved hands	Cooked ham	Yes	Pérez-Rodríguez et al., 2006

In these studies factors enhancing the amount of *L. monocytogenes* being transferred were a high amount of bacteria, or biofilm; stainless steel surface compared to polyethylene surface; hydration of biofilm and higher force of the cutting blade. Additionally food type and the presence of non-*Listeria* species on the surface had a clear impact on transfer (Midelet and Carpentier, 2002; Lin et al., 2006; Vorst et al. 2006a; Rodríguez and McLandsborough, 2007). Ivanek et al. (2004) and Schaffner (2004) presented mathematical models describing *L. monocytogenes* cross-contamination in food-processing plants.

1.4 Prevention of *L. monocytogenes* in food processes

In the following, aspects on prevention of *L. monocytogenes* in food processes are presented with special focus on disinfection.

1.4.1 General aspects in prevention

As *L. monocytogenes* is widely distributed in nature and survives in various environmental conditions, total elimination of this bacterium from most food processes is impractical or even impossible. However, it is possible to reduce and control the level of this bacterium and minimize the risk to public health. By using a variety of physico-chemical factors, singly or in combinations, effective control of growth and survival of *L. monocytogenes* can be achieved. It has been suggested that the cause of *L. monocytogenes* contamination in products is due more to contamination from the processing environment and equipment than from the raw materials (Bell and Kyriakides, 2004). Strict temperature control and storage time limitations in products supporting the growth, and absence or lowest possible initial level of the bacterium, are needed to minimize the occurrence and amount of this bacterium in food products (FAO/WHO, 2004). This pinpoints the importance of a high level of equipment hygiene.

Tompkin (2002) presented six strategies for controlling *L. monocytogenes* in food plants. These included (I) prevention of the establishment and growth of *Listeria* in niches or other sites that can lead to the contamination of RTE foods; (II) implementation of a sampling program that can assess whether the environment is under control; (III) as rapid and effective a response as possible to each positive product contact sample; (IV) verification by follow-up sampling that the source has been detected and corrected; (V) a short-term assessment of last samplings to detect problems and trends and (VI) a longer-term assessment.

Examples of successful eradication means of *L. monocytogenes* have been published. In a study by Autio et al. (1999) of *L. monocytogenes* contamination in a cold-smoked rainbow processing plant, an *L. monocytogenes* eradication program consisting of the use of hot steam, hot air and hot water was successfully implemented. None of the control samples taken five months after eradication contained *L. monocytogenes*. In a study by Samelis et al. (1998),

cooked ham products were contaminated with *L. monocytogenes* during tumbling step, but the product was absent from vacuum packed product provided that heating to a core temperature of 70 °C occurred and recontamination during slicing and packing was prevented. *L. monocytogenes* has been found to contaminate products from the body of the meat slicer switches. In this case a simple solution, rubber switch covers, solved the problem (Lelieveld et al., 2003).

1.4.2 Efficiency of disinfectants on L. monocytogenes

L. monocytogenes, as in general for gram-positive bacteria, is more sensitive to disinfectants than most gram-negative bacteria. Gram-positive bacteria lack the outer surface layer of gram-negative bacteria, which restricts the entry of various antimicrobial substances (McDonnell and Russell, 1999). A summary of the effects of a variety of disinfecting compounds on *L. monocytogenes* in different test conditions is presented in Table 7. Due to many varying factors in the disinfect tests the efficacy against *L. monocytogenes* is difficult to compare. Furthermore, commercial products containing a variety of compounds are also difficult to compare. As can be concluded from the studies presented in Table 7 and from studies made with commercial products (Sallam and Donnelly, 1992; Aarnisalo et al., 2000), many disinfecting or sanitizing agents commonly used in the food industry, such as quaternary ammonium compounds (QACs), chlorine and iodofors, are effective against *L. monocytogenes* cells in suspension.

The microbicidal effect of quaternary ammonium compounds is based on membrane damage in the phospholipid bilayers. Halogens e.g. chlorine inhibit DNA-synthesis and cause oxidation of thiol groups in proteins and enzymes. Oxidation is also caused by peroxygens. The mode of action of alcohols is not well known, but most probably they cause membrane damage and denaturation of proteins (McDonnell and Russell, 1999).

The surface attachment and biofilm formation of *L. monocytogenes* (Mustapha and Liewen, 1989; Frank and Koffi, 1990; Dhaliwal et al., 1992; Wirtanen and Mattila-Sandholm, 1992b; Mosteller and Bishop, 1993) as well as the presence and type of organic material (El-Kest and Marth, 1988a; Best et al., 1990; Wirtanen and Mattila-Sandholm, 1992a; van de Weyer et al., 1993; Aarnisalo et al., 2000) affects the disinfectant efficacy and a thorough cleaning of the

surfaces should be performed before disinfection (Krysinski et al., 1992). Other factors affecting the disinfectant efficacy against *L. monocytogenes* include the concentration, duration, pH and temperature of the use solution (El-Kest and Marth, 1988b; Tuncan, 1993). Furthermore, variation between *L. monocytogenes* strains in resistance against disinfectants has been reported (Earnshaw and Lawrence, 1998; Teodorovic et al., 2000; Lundén et al., 2003).

Only a few publications are available on disinfectant efficacy against L. monocytogenes at low temperatures (Orth and Mrozek, 1990; Tuncan, 1993), although L. monocytogenes is problematic especially in cold areas of food processing, e.g. in the dairy and meat industry (Lundén, 2004; Wirtanen and Salo, 2004). Coolers and freezers are potential habitats for this bacterium (Gravani, 1999), and efficient disinfection of these premises must be performed. In the study of Tuncan (1993) the efficacy of disinfectants containing quaternary ammonium compounds and iodofors in low concentrations (50 ppm) was clearly reduced when the temperature was reduced from 25 °C to 2 °C. Cold temperature did not affect the efficacy of chlorine (25-200 ppm). However, increased effectiveness of chlorine due to an increase in temperature has also been reported (El-Kest and Marth, 1988b; Orth and Mrozek, 1990). Additional factors reported to affect the efficacy include the quality of used water (Marriott, 1999), the age of the bacterial cells (El-Kest and Marth, 1988a; Lee and Frank, 1991b), the time that biofilm has grown on a surface (Shin-Ho-Lee and Frank, 1991), the surface material (Krysinski et al., 1992; Mosteller and Bishop, 1993; Ronner and Wong, 1993) as well as the nutrient level and growth temperature of the L. monocytogenes cells (Lee and Frank, 1991a).

In places which are difficult to reach *L. monocytogenes* may encounter suboptimal concentrations of disinfectants. Such places are common in unhygienic food processing equipment. It is possible that the resistance of the organism towards the agent increases through adaptive responses (Aase et al., 2000). This can also occur through cross-adaptive response to different disinfectants (McDonnell and Russell, 1999; Lundén et al., 2003). Adaptive response is a characteristic of *L. monocytogenes*, which partly explains its ability to persist in food processing equipment (Lundén, 2004; Møretrø and Langsrud, 2004).

Disinfecting agent	Use-concentration		Tes	t conditi	SUC		Efficacy ^a		Reference
	(mdd)	t (min)	T (°C)	Clean	Soiled (soil type and load)	Suspension	Surface ^b		
Ethanol	70% vol/vol	-	20	TSB		‡	+		3est et al., 1990
					whole human serum pasteurized milk (2% fat)	‡ ‡	• +		
Sodium hypochlorite	10 60 (AV CI 6)	-	00	TSR		^b UN/++	++/-		Aest et al 1000
			Ì		whole human serum	+++++++++++++++++++++++++++++++++++++++	-/-		
					pasteurized milk (2% fat)	++ / ++	+/+		
	200 (AvCl)	10	20	TSB		QN	- / - c/ - f		Krysinski et al., 1992
	200 (AvCl)	0.5	20	TSB		ŊŊ	+ (4h) ^g / - (8d) ^g		Shin-Ho-Lee and Frank, 1991
	200 (AvCl)	0.5	25			‡	ND		Mosteller and Bishop,
	200 (AvCl)	0.5	25		steamed milk (2% fat)	QN	+ h, i		
	50, 100, 200, 400, 800 (AvCl)	1, 2, 5	20	TSB		+/++(≥200 ppm)	+/++(≥800, ≥400, ≥800ppm) ^j +/++(≥800 ppm) ^k	(11) ^g (11) ^g	Mustapha and Liewen, 198
	50, 100, 200, 400, 800 (AvCl)	1, 2, 5	20	TSB		+/++ (≥200 ppm)	$+/++(\geq 400, \geq 400, \geq 400 \text{ ppm})^{1}$ +/++($\geq 800 > 400 > 200 \text{ rnm}^{k}$	(24h) ^g (24h) ^g	
	25, 50, 100, 200 (AvCl)	0.5	2; 7; 15.5; 25	saline		++ / ++ / ++ / ++	ND		Funcan, 1993
<u>Iodofor</u>	00 (1-1-1-1)	-	ç	ЦЪТ.		-			0001 1
	00 (UULALAULE 12)	-	70	gei	سابعه مصيبا واصلينا	F			JCSI CI AI., 1990
					pasteurized milk (2% fat)	• ‡			
	25	10	20	TSB		ND	-/- ^e /- ^f		Krysinski et al., 1992
	25	0.5	25			+	ND		Mosteller and Bishop, 1993
	25	0.5	25		steamed milk (2% fat)	ND	+ h, i		
	25, 50	0.5	2	saline		+/+	ND		Funcan, 1993
	25, 50	0.5	7	saline		++ / +	ND		
	75 50	20	15 5.75	colino.		++/++			

Disinfecting agent	Use-concentration		Tes	t conditie	SUC	Ð	fficacy ^a	Reference
	(mdd)	t (min)	T (°C)	Clean	Soiled (soil type and load)	Suspension	Surface ^b	
Quaternary ammonium		,	1					
compound	400, 500	-	20	TSB		UN / ++	++/+	Best et al., 1990
					whole human serum	+/+	- / -	
	200 (pH 4.4 or 6.4)	10	20	TSB		‡ Q	+ + / - e/ - f	Krvsinski et al 1992
	200	0.5	25			‡	QN	Mosteller and Bishop,
	200	0.5	25		steamed milk (2% fat)	QN	- h / + i	6661
	50, 100, 200, 400, 800	1, 2, 5	20	TSB		‡	‡	Mustapha and Liewer
	25, 50, 100, 200	0.5	7	saline		++ / ++ / + / +	QN	1989 Tuncan, 1993
	25, 50, 100, 200	0.5	7; 15.5			++ / ++ / ++ / +	ND	×
	25, 50, 100, 200	0.5	25			++ / ++ / ++ / ++	ND	
Peracetic acid		\$	ę	40£		Ę	یں 19 	
	200	0.5	25 25	gei		N ‡	- / + / + GN	Mosteller and Bishop,
	200	0.5	25		steamed milk (2% fat)	ND	+ h / _ i	1993
Acid anionic	200	0.5	25			ŧ	Ę	Mosteller and Bishon
	200	0.5	25		steamed milk (2% fat)	ŊŊ	+ ^h , i	1993

 - = no efficacy (<1 log 	
a ++ = good efficacy (\ge log reduction in suspension, \ge 4 log reduction on surface), + = poor efficacy (<5 log reduction in suspension, <4 log reduction on surface),	^b stainless steel, ^c available chlorine, ^d not detected, ^e polyester, ^f polyester/polyurethane, ^g attachment time, ^h buna-N-rubber, ⁱ Teflon [®] , ^j smooth, ^k pitted

1.5 Microbiological risk assessment (MRA) in the food industry

1.5.1 Background and components of MRA

Microbiological risk assessment was launched after the implementation of the Sanitary and Phytosanitary Agreement by the World Trade Organization (WTO) (WTO, 1994) in 1995 in order to assist in fulfilling and clarifying global trade regulations, and has been used by national and international authorities in risk management decision making. Codex Alimentarius (1999)defines microbiological risk analysis to consist of three components: risk assessment, risk management and risk communication. The microbiological risk assessment process should have a sound scientific basis, be transparent, and be conducted using a structured approach including hazard identification, exposure assessment, hazard characterization and risk characterization. It should explicitly consider the dynamics of microbiological growth, survival and death in foods and the interaction between human and agent following consumption.

The purpose of the hazard identification is to identify the micro-organisms or the microbiological toxins of concern with food. Exposure assessment includes an assessment of the extent of human exposure by considering the frequency and levels of contamination of foods over time and at the time of consumption. In the hazard characterization step, a qualitative or quantitative description of the severity and duration of possible adverse effects from consumption of a micro-organism or its toxin is provided. Ideally, a dose-response relationship should be established but in the absence of a known dose-response relationship, expert opinions can be used. Risk characterization combines information obtained from hazard identification, hazard characterization and exposure assessment to obtain a risk estimate, which is a qualitative or quantitative estimate of the likelihood and severity of the adverse effects which could occur in a given population (Codex Alimentarius, 1999).

1.5.2 Microbiological risk assessment of L. monocytogenes

FAO/WHO (2004) has conducted a risk assessment of *L. monocytogenes* in four categories of RTE foods, namely pasteurized milk, ice cream, fermented meat and cold-smoked fish. It was concluded that the food matrix, virulence of the strain and susceptibility of the consumer were all important factors for the probability of listeriosis. The models developed predicted that a high number of *L. monocytogenes* is needed for illness and that the greatest impact on reducing rates of listeriosis would be obtained by reducing high levels of contamination. In foods where growth of *L. monocytogenes* is supported, better temperature control and limitation of storage times would be beneficial. Other risk assessments of *L. monocytogenes* have been performed with the following products: soft cheese from raw milk (Farber et al., 1996; Bemrah et al., 1998; Sanaa et al., 2004), Hispanic-style cheese (FAO/WHO, 2004), Swiss Emmental cheese (Aebi et al., 2003), smoked fish (Buchanan et al., 1997; Lindqvist and Westöö, 2000), Deli meats (FSIS, 2003), RTE products (FDA/USDA, 2003) and butter (Maijala et al., 2001).

For *L. monocytogenes*, dose-response relationship models are available from some studies and have been summarized in the FAO/WHO document on risk assessment of *L. monocytogenes* in RTE foods (FAO/WHO, 2004). A variety of dose-response models based on epidemiological data (Buchanan et al., 1997), expert elicitations (Farber et al., 1996) and data derived from surrogate pathogens or animals (Haas et al., 1999), are available for risk assessment of *L. monocytogenes* (FAO/WHO, 2004). These generally include the use of the exponential model (Rose et al., 1991), but e.g. the Beta-Poisson (Haas, 1983) and Weibull-Gamma models (Todd and Harwig, 1996) are also used.

1.5.3 Use of predictive models in MRA and risk management

Predictive microbiology can be used to estimate changes in bacterial numbers, allowing the exposure of an individual to a pathogen to be assessed. Without predictive models this has been difficult to carry out in microbiological risk assessment (Walls and Scott, 1997). Predictive microbiological growth models can be divided into primary, secondary and tertiary models. Models that describe the population growth curve are primary models (Whiting and Buchanan, 1993).

This description requires definition of the initial number of cells, the lag time, the rate of growth and the maximum population density. Models describing inactivation of micro-organisms have also been developed (van Gerwen and Zwietering, 1998). Secondary models describe the influence of environmental conditions on the parameters of the primary model, or the primary characteristics of the population and environment interactions, e.g. lag time of the microbe and exponential growth rate. Tertiary models are computer software interfaces that enable rapid predictions of microbiological growth from values selected by the user (Ross and McMeekin, 2003).

Predictive models can be used as tools for process development and optimization in predicting the bacterial levels in the final products and the effect of cross-contamination, as described e.g. for *L. monocytogenes* by Rasmussen et al. (2002) and Ivanek et al. (2004). Many models are also available in computer programs such as Pathogen Modeling Program (PMP, <u>http://www.arserrc.gov/mfs/pathogen.html</u>), and Seafood Spoilage and Safety Predictor (<u>http://www.difres.dk/micro/sssp/</u>). Data underlying many predictive models can be found in ComBase (<u>http://www.ifr.ac.uk/combase/</u>). The predictive models have limitations, which should be recognized in order to avoid unrealistic scenarios. Such limitations include uncertainty, which is the expression of knowledge gaps, and variability which describes the heterogeneity of the quantities and characteristics modelled (Ross and McMeekin, 2003).

1.5.4 Use of MRA and predictive models at the food plant level

Although risk assessment has been used mainly on a national and international level, it is a beneficial approach for food companies in product and process development and optimization, as an extension of or in validation of an HACCP-plan (Notermans and Mead, 1996; van Schothorst, 1997; Serra et al., 1999; Hoornstra et al., 2001; Hoornstra and Notermans, 2001). In risk assessments for governments use is mainly made of epidemiological data, whereas for food companies product and process information is generally used. In most HACCP-systems a qualitative approach is used. Quantitative risk assessment is still challenging for producers, but it could be used to quantify the effect of control measures, to estimate the occurrence of contaminants in the end products and in deriving and validating control measures and critical limits at Critical Control

Points (CCPs). Either worst-case or what-if scenarios, or a statistical approach using probability distributions can be applied (Hoornstra et al., 2001). At the plant level, a compact and simple way of performing the risk assessment is needed. Van Gerwen et al. (2000) presented a general method for stepwise quantitative risk assessment of food products and their production processes. Computer programs for performing a risk assessment have recently been launched (Ross and Sumner, 2002; Sumner and Ross, 2002; Tuominen et al., 2003; McMeekin et al., 2006), e.g. a practical semiquantitative hygiene risk assessment model HYGRAM[®] by Tuominen et al. (2003). This model includes a hazard module for assessing the risk of *L. monocytogenes*. The principles of risk assessment underlying the programs should also be familiar to their users. Predictive microbiology models provide a scientific basis to support key aspects of HACCP and quantitative microbiological risk assessment (Kleer and Hildebrandt, 2001; McMeekin et al., 2006).

2. Aims of the study

The aim of this thesis was to identify deficiencies in and to improve means of equipment and process hygiene in the food industry and to develop risk assessment procedures for prevention of *L. monocytogenes* in food plants. Based on the results, suggestions for improved manufacturing and risk management practices for assuring end product safety are given. Specific aims were:

- 1. To identify and evaluate food processing equipment and the hygienic practices of maintenance personnel in the food industry in order to identify aspects that can negatively affect equipment hygiene and enhance the occurrence of *L. monocytogenes* in equipment (Paper I).
- 2. To compare the discriminatory power of automated ribotyping to that of PFGE in distinguishing *L. monocytogenes* isolates and to determine the suitability of the methods for tracing contamination sources in food processes (Paper II).
- 3. To enhance means of *L. monocytogenes* decontamination in equipment by evaluating commercial lubricants and disinfectants used in the food processing equipment; by determining the survival, growth and transfer of *L. monocytogenes* in lubricants (Paper III); and by investigating the susceptibility of the bacterium to disinfectants in cold conditions (Paper IV).
- 4. To develop risk assessment procedures by using predictive modelling to investigate transfer of *L. monocytogenes* from equipment to product during slicing to assess recontamination (Paper V); and by performing a production plant level risk assessment of *L. monocytogenes* for one food product (Paper VI).

3. Materials and methods

The mail-surveys performed in equipment and maintenance hygiene studies are described (Paper I). The methods used for sampling and analyses of *L. monocytogenes* (Papers I–VI) are summarized and methods used in susceptibility (Papers III and IV), transfer (Paper V) and risk assessment (Paper VI) studies, as well as the mathematical methods used in the studies, are also described. Cold-salted (i.e. 'gravad') salmon (Paper V) and raw marinated broiler legs (Paper VI) were chosen as example foods in laboratory experiments because *L. monocytogenes* is often detected in these products (see Table 5).

3.1 Questionnaires on equipment hygiene and hygienic working practices of maintenance personnel (Paper I)

Two mail surveys to Finnish food companies were sent in spring 2002 to 1) identify and evaluate equipment causing hygiene problems in food processing and 2) to study the hygienic practices of maintenance personnel in the food industry in order to identify aspects that can have a negative effect on equipment hygiene. The respondents were able to answer anonymously. The companies were chosen from the company registers of the Finnish Food and Drink Industries' Federation and VTT.

The questionnaire on equipment hygiene was sent to 184 food companies. An employee responsible for equipment hygiene was asked to answer the survey. The questionnaire on hygienic working practices of maintenance personnel was sent to the quality managers of 106 food companies, who were asked to distribute it in addition to themselves to maintenance personnel (330), food-handlers (118), and cleaning personnel (224), i.e. a total of 778 questionnaires. More detailed descriptions of the surveys can be found in Paper I.

3.2 Sampling, detection and identification of *L. monocytogenes*

3.2.1 L. monocytogenes strains used in laboratory experiments

The *L. monocytogenes* strains used in laboratory experiments (Papers III and IV) were obtained from the VTT culture collection (except strain F2365 in Paper V, which was obtained from the United States Department of Agriculture) and maintained in 5% glycerol at -70 °C. For each study the strains were chosen mainly on the basis of their site of isolation and the food sector they represented. The strains used are presented in Table 8. For comparing the discriminating ability of automated ribotyping to that of pulsed-field gel electrophoresis (PFGE) in distinguishing *L. monocytogenes* isolates (Paper II), a total of 486 *L. monocytogenes* isolates originating from 17 Finnish food processing plants were collected from 1997 to 1999. For further typing with *Pvu*II enzyme as well as with PFGE, a set of 121 isolates was selected from these isolates, representing all the *Eco*RI ribotypes generated and 16 food plants.

Strain	Origin	Serotype	Papers
I (VTT E-981041)	Meat plant, product	1/2	IV
II (VTT E-991508)	Dairy, cheese machine	1/2	IV
III	Meat plant, freezer	4b	III, IV
IV (VTT E-021893)	Meat plant, freezer	1/2	IV
V	Meat plant, product	4b	IV
VI (VTT E-991599)	Meat plant, product	1/2	IV
VII (VTT E-991513)	Fish plant, fish net	1/2	IV
VIII	Meat plant, conveyer	1/2	IV
IX (VTT E-981012)	Dairy, raw milk	1/2	III, IV
X (VTT E-981045)	Meat plant, product	1/2	IV
XI (VTT E-991205)	Dairy, butter	3a	III
F2365	Isolate from the 1985 Mexican-style soft cheese outbreak (Linnan et al., 1988), rifampicin-resistant mutant	4b	V

Table 8. Strains of Listeria monocytogenes used, their origin, and serotypes.

3.2.2 Analyses of tools, work environment and protective clothing of maintenance personnel (Paper I)

In addition to the questionnaires (see 3.1), the working practices of maintenance personnel were studied at four food companies (a meat company, a poultry company, a dairy and a bakery) during two normal work shifts by taking microbiological samples. For analysis of *L. monocytogenes* a total of 71 samples from three food plants were taken with moisturised gauze pads kept in 10 ml of a peptone saline solution (Maximal Recovery Diluent, Lab M, Amersham, Bury, UK). The isolation and detection were carried out according to the ISO 11290-1 method (ISO, 1996) with the following modifications: only Oxford agar (Oxoid, Hampshire, UK) was used as the selective agar and the preliminary identification was carried out using API Listeria strips (bioMérieux, Marcy-l'Etoile, France) in accordance with the manufacturer's instructions.

Additionally, samples for analysing total aerobic bacteria were taken with commercial PetrifilmTM Count Plates (3M Health Care, St. Paul, MN, USA) and for *Enterobacteriaceae* with 3M PetrifilmTM Enterobacteriaceae Count Plates from four food plants. Altogether, 95 and 96 samples were taken and incubated at 30 °C for 3 d and at 37 °C for 2 d, respectively. After this the colonies were counted.

3.2.3 Analyses of cold-salted salmon slices in a transfer study (Paper V)

For analyzing the amount of *L. monocytogenes* transferred during slicing of cold-salted salmon from inoculated blade to product (see 3.5), the slices (i.e. samples) were transferred directly to sterile filter (280 μ m mesh) stomacher bags (Spiral Biotech, Norwood MA, USA) and weighed for more accurate enumeration. Peptone water (0.1% wt/wt) was added to the sample at a 5:1 ratio (wt/wt) and the mixture was processed in a stomacher (Model Bag Mixer 400, Interscience Inc., Weymouth, MA, USA) at room temperature for 30 s. The swabs used for surface sampling were placed individually into 10 ml of peptone water, mixed in a vortex mixer for 30 s, and then 10-fold serial dilutions (100 μ l from each tube) were plated on two plates of Modified Oxford (MOX) -agar

containing 0.1% rifampicin (Sigma Aldrich, St Louis, MO, US). The plates were incubated at 37 $^{\circ}$ C for 2 d and the colonies were counted.

3.2.4 Analyses of broiler legs in a risk assessment study (Paper VI)

For analysing the prevalence and numbers of *L. monocytogenes* in raw marinated broiler legs for the risk assessment study (see 3.6.5), a total of 186 packages of raw marinated broiler legs were purchased from 41 retail stores in the Helsinki metropolitan area during one year in 2002–2003. The packages were transferred to the laboratory in an insulated box, and were stored at 6 °C or 10 °C, and analysed at the end of their shelf life. To quantify *L. monocytogenes*, one leg from each package was transferred to a sterile bag and weighed, and half of the sample weight of buffered peptone-water was added. The broiler leg was hand massaged for 3 min, followed by a pre-incubation for 1 h at room temperature. The rinse diluent was used for enumeration and isolation conducted according to the ISO 11290-2 method (ISO, 1998) and the detection of *L. monocytogenes* after heating experiments according to the ISO 11290-1 method (ISO, 1996), with some modifications as described in Paper VI.

Information on levels of *L. monocytogenes* at the point of consumption is needed when performing risk assessment and therefore a small scale heating experiment was performed. For investigating the temperatures normally used by consumers cooking broiler legs, cooking practices of 20 consumers, mainly students and young adults, were investigated using a temperature logger (DataSquirrel, Eltek Limited, UK). Consumers were given a package of broiler legs, corresponding to those investigated in the microbiological survey, and asked to cook them in their home kitchen oven as they normally would. No instructions were given except on the correct use of the temperature logger, which measured both the oven's air temperature and the meat temperature at the thickest portion of the leg. They were also asked to fill in a small questionnaire concerning their cooking practices.

The thermal inactivation of *L. monocytogenes* from naturally contaminated broiler legs was investigated in the laboratory from 21 samples kept at 10 °C. The packages (samples) purchased from retail shops contained 3–4 broiler legs in a modified atmosphere package. The numbers of *L. monocytogenes* was investigated from one leg and two legs were cooked one at 132 °C and the other

at 175 °C, for 50 min. The former temperature was chosen as it was the lowest average temperature detected in consumer ovens during cooking. The other temperature was the temperature recommended by the producers. The cooking time was the shortest recommended by the producers. After heating, the samples were cooled and stored at 10 °C until the next day when numbers of *L. monocytogenes* were determined. A more detailed description of the experiments is presented in Paper VI.

3.2.5 Typing of L. monocytogenes (Paper II)

The discriminatory power and usability of automated ribotyping was compared with traditionally used PFGE for distinguishing *L. monocytogenes* strains isolated from food processing plants. Additionally the strains were serotyped. The isolates were ribotyped using the RiboPrinter[®] System (DuPont QualiconTM, Inc., Wilmington, DE, USA) as described by Bruce (1996). The restriction enzymes used were *Eco*RI (Qualicon) and *Pvu*II (Qualicon). Similarity values were calculated using the software provided by Qualicon (version 11.2 (c) 1999).

In situ DNA isolation and PFGE were performed as described by Autio et al. (1999). The restriction enzymes used were *AscI* and *SmaI* (New England Biolabs, Beverly, Mass., USA). The serotyping was carried out by the agglutination method using Denka Seiken's *L. monocytogenes* serotyping antisera (Denka Seiken, Tokyo, Japan) according to the manufacturer's instructions.

3.3 Analysing survival and transfer of *L. monocytogenes* in lubricants (Paper III)

3.3.1 Survival of *L. monocytogenes* in lubricants

The survival of *L. monocytogenes* in lubricants used in food processing equipment was investigated in 11 different types of lubricants used in food processing plants (Table 9). A mixture of 0.1% bovine albumin and 0.1% potato starch was added as soil. Culturing of the *L. monocytogenes* strains (Table 8) is presented in detail in Paper III. The inoculated tubes were incubated both at room (20 °C) and refrigerated (5 °C) temperatures. The samples (0.5 g) were

taken in triplicate after 0.5 h (control sample), 4 h, 24 h, 3 d and 14 d and pipetted to 4.5 ml of neutralization solution (Aarnisalo et al., 2000) and the solution was allowed to stand for 5 min before culturing. To enhance the emulsification of the lubricants, a dilution series was prepared in pre-warmed (35 °C) 0.85% Tween-saline solution, plated on duplicate plates of Oxford agar (Oxoid Ltd., UK) and incubated at 30 °C for 3 d. The microbicidal effect (*ME*) of the lubricants was calculated as described in 3.6.1.

3.3.2 Transfer from lubricants to stainless steel surfaces and vice versa

For investigating transfer from lubricants to stainless steel surfaces the first seven lubricants (A–G) given in Table 9 were soiled and inoculated with strains III and XI (VTT E-991205). The lubricant (2 g) was transferred to a sterile Petri dish containing a filter paper (Whatman, qualitative no. 2, Maidstone, UK) and spread to cover the whole paper. Sterile stainless steel discs were placed on top of the filter paper and after 0.5 h,1 h, 4 h and 24 h three discs were analyzed as replicate samples. They were transferred to test tubes containing 5 ml of neutralization solution. The lubricant-contaminated stainless steel disc was mixed with the inactivation solution in a test tube mixer and the disc was further rubbed with a cotton swab for 30 s. The neutralization solution was allowed to stand for 5 min before culturing. After the same time periods (0.5 h, 1 h, 4 h and 24 h), samples were taken from the lubricant in order to detect changes in the *L. monocytogenes* concentration in the lubricant during the 24 h incubation period.

Lubricant	Composition	Usage
A^1	Synthetic hydrocarbon (70–80%), hydro treated polymer (10–20%), antioxidant (0.5–2%), fumed silica (7–10%), thermal stabilizer (3–5%), polyglycol (1–2%), polytetrafluorethylene (Teflon) (0.5–2%), additives (0.25–1%)	Food industry lubricant (grease, USDA H1)
B^2	Silicone, non-ionic surface active compounds (<5%), preservatives	Conveyer belt lubricant (3 ml/L aqueous solution of lubricant in use)
C^2	Crude oil (30-60%), butane (20-50%), propane (5-30%)	Chain lubricant
\mathbf{D}^1	Rapeseed oil (100%)	Cooking oil
E^2	Mineral oil containing a mixture of mainly saturated hydrocarbons with C15–C50 (>90%), additives	Hydraulic oil
F^2	"Dry" (not diluted with water), polyhydric alcohols and a small amount of silicone emulsion	Conveyer belt lubricant
G^1	White oil (>97%), dialkyl-dimethyl-aluminum silicate, additives (against corrosion, wear, pressure and oxidation)	Food industry lubricant (grease, USDA H1)
H^2	Mineral oil containing a mixture of mainly saturated hydrocarbons with C15–C50 (>90%), additives	Gear oil
I^2	Mineral oil containing a mixture of mainly hydrocarbons with C12–C50 (>90%), lithium thickener, additives	Multi-purpose grease for vehicles
\mathbf{J}^1	White oil, Al-salt of stearic acid (5–15%), Al-salt of benzoic acid $(1-5\%)$	Aluminium complex grease (USDA H1)
K ²	Hydrodesulfurized light dearomatized naphtha (petroleum) (30–60%), hydrocarbon propellant (10–30%), additives	Multi-purpose grease

Table 9. Lubricants, composition and usage as given by the manufacturers.

¹ food grade ² non food grade

For investigating transfer of bacteria from surfaces into lubricants, growth broth (0.1 ml) containing $10^8 \text{ CFU/ml} L$. *monocytogenes* was pipetted onto the surface of sterile stainless steel discs (#AISI 304, 2B, 10.5 mm in diameter, Happoteräs Oy, Helsinki, Finland). The discs were left to dry for 1.5 h at 30 °C, after which they were further dried for 10 min with compressed air. The discs were transferred to 10 g of soiled lubricant. After 1 h, 4 h and 24 h, three discs were removed from the test tubes and the tubes were incubated at room (20 °C) and refrigerated (5 °C) temperatures. The soiled lubricants were mixed and three replicate samples from each tube were plated as described in 3.3.1.

3.4 Susceptibility of *L. monocytogenes* to disinfectants (Paper IV)

The efficacy of commercially available disinfectants commonly used in the food industry against *L. monocytogenes* at cold temperature (+5 °C) was investigated using both suspension and surface methods.

3.4.1 Suspension method

The efficacy of eight disinfectants (Table 10) was studied in both clean and soiled solutions with ten strains I–X (Table 8) using a method slightly modified from Aarnisalo et al. (2000). A more detailed description of the test is given in Paper IV. Microbicidal effect (ME) was calculated as described in 3.6.1. The disinfectant was considered to be effective if it reduced the amount of vegetative bacterial cells by 5 log CFU-units.

3.4.2 Surface method

The surface method used was modified from Charaf et al. (1999) using 5 strains (Table 8) and disinfectants presented in Table 10. The surfaces used were stainless steel discs (AISI 304, 2B, 10.5 mm in diameter, Happoteräs Oy, Helsinki, Finland) and glass bead blasted PE-discs (10.5 mm in diameter, Vulganus Oy, Nastola, Finland). The culturing, inoculation and disinfection efficacy methods are described in detail in Paper IV. Microbicidal effect (ME) was calculated as described in 3.6.1. The disinfectant was considered to be effective if it reduced the amount of vegetative bacterial cells by 4 log CFU-units.

Table 10. The composition, purpose of use and recommended in-use concentrations of disinfectants.

Agent	Composition	Purpose of use	Recommended in- use concentrations (%)
Α	Peracetic acid (<5%), hydrogen peroxide (15–30%), acetic acid (5–15%), phosphonic acid (<1%)	Disinfectant for closed processes	0.05–3
В	Peracetic acid (<5%), hydrogen peroxide (5–15%), acetic acid (5–15%), phosphonic acid (<5%), non-ionic tensides (<5%)	Disinfectant for open processes	1–3
С	Peracetic acid (<5%), hydrogen peroxide (5–15%), acetic acid (15–30%), phosphonic acid (<5%), anionic tensides (5–15%)	Disinfectant for closed processes	0.05-1
D	Ethanol (<70%)	Disinfectant for open processes	100
Е	Sodium hypochlorite (>60%, active chlorine 13%), sodium hydroxide (<5%)	Disinfectant for open processes	0.05–2
F	Alkyl dimethyl benzyl ammonium chloride (>30%), synthetic tensides	Disinfectant for closed processes	0.1–0.5
G	Isopropanol (15–30%), 1-propanol (>30%)		100
Н	Peracetic acid (5–15%), hydrogen peroxide (15–30%), acetic acid (5–15%)	Disinfectant for closed processes	0.1–4

3.5 Analysing transfer of *L. monocytogenes* during slicing of cold-salted salmon (Paper V)

Fresh Atlantic salmon (*Salmo salar*) fillets were stored at 0 °C for up to experimentation time (no longer than one week). The fillets were salted as described in detail in Paper V. The thickness of the slices was set at 3.5 mm and slicing was performed manually. After each trial, the slicer and blade were cleaned and disinfected. All trials were repeated three times on different days.

The slicer used in this study was a delicatessen slicer (Globe 3975 Variable Speed Automatic Slicer, Globe Food Equipment Co., Dayton, OH, USA) made of #304 stainless steel, as also was the blade. The roughness (A'/A ratio, which is the scanned rough surface area/unit surface area) and sharpness of the blade

were measured at the beginning and the end of the tests by reflection confocal microscopy as described by Flores et al. (2006). The blade was contaminated with *L. monocytogenes* resulting in approximately 8, 5 or 3 log CFU/blade. The effect of slicing temperature was studied at room temperature, 10 ± 0.5 °C, and 0 ± 0.2 °C. The slicer and blade were pre-cooled to the experimental temperature before inoculation. For experiments, the first 11 slices and then every second slice to the 39th slice were collected. Surface samples (from approximately 10 cm² area each) were collected from the slicer blade, holding plate and blade safety guard using sterile cotton-tip swabs during slicing.

Transfer from inoculated salmon fillet to slicing machine and to slices of uninoculated fillets was studied by surface-inoculating the salmon fillet to contain approximately 8 log CFU of *L. monocytogenes*. The trials were made at room temperature. After slicing the inoculated fillet, an uninoculated fillet was sliced into 39 slices and the slices were analyzed. Surface samples as described above were collected.

3.6 Mathematical methods

3.6.1 Microbicidal effect (Papers III, IV)

The microbicidal effect (ME) of lubricants and disinfectants was calculated according to the following Equation (1):

$$ME = \log N_c - \log N_d, \tag{Eq. 1}$$

where N_c is CFU/ml or CFU/cm² for the control and N_d is CFU/ml or CFU/cm² for the treatment.

3.6.2 Discrimination index (Paper II)

The discrimination power of the automated ribotyping and PFGE was determined by calculating the discrimination index (*DI*) using the following Equation (2):

$$DI = 1 - \frac{1}{N(N-1)} \sum_{j=1}^{S} n_j (n_j - 1)$$
(Eq. 2)

where N is the total number of strains in the sample population, s is the total number of types described and n_j is the number of strains belonging to the *j*th type (Hunter and Gaston, 1988).

3.6.3 Statistical analyses (Papers I, III, V)

In the study on survival and growth of *L. monocytogenes* in lubricants (Paper III), repeated measures analysis of variance was used to analyze whether there were significant differences in reduction in the amount of *L. monocytogenes* due to change in time, lubricant, temperature, purity of lubricant and bacterial strain. Differences within the lubricants and different strains were further analyzed by the multiple comparisons Tukey-test.

The significance of different experiment conditions on transfer of *L. monocytogenes* to slices of cold-salted salmon (Paper V) were analyzed with a general linear model univariate analysis. Differences in numbers of aerobic bacteria on maintenance personnel tools, protective clothing and in the work environment (see 3.2.2) were calculated using the non-parametric Mann-Whitney test.

All statistical analyses were conducted using SPSS for Windows v. 12.0.1. (Chicago, USA). The level of significance used was p < 0.05.

3.6.4 Predictive modelling (Paper V)

TableCurve 2D Version 5.01 (SYSTAT Software Inc., Richmond, CA, USA) was used in the study on transfer of *L. monocytogenes* in order to select an empirical model to best fit the experimental data, based on the simplicity, applications (predictions vs. time in slicing – convergence and no singularity in long time prediction), fitted coefficients with standard error, *t*-test results, P > |t|, r^2 -value and *F*-value.

3.6.5 Producer level quantitative risk assessment (Paper VI)

An example of a robust risk assessment for the single plant level using worstcase and average point estimates was produced. The steps of microbiological risk assessment are hazard identification, exposure assessment, hazard characterization and risk characterization (see 1.5). For exposure assessment purpose, the prevalence and numbers of *L. monocytogenes* in marinated broiler legs were investigated and a laboratory scale heating study was performed to estimate the numbers of *L. monocytogenes* at the point of consumption (see 3.2.4). A scheme/flowchart of the factors and probabilities needed for producing the risk estimate was drawn (Fig. 1, Paper VI).

Output estimates of *L. monocytogenes* -positive samples were then compared to the data obtained in the cooking studies (see 3.2.4). Since the cooking data were very limited, the assessed probabilities of cooking time (min) and oven temperature (°C) combinations (t,T-values) were sensitive to possible outliers in the data. Therefore, for assessing the probability of the highest risk cooking combinations, a simple simulation which could be performed using e.g. Excel was performed. Natural logarithms of t (time, min) and T (temperature, °C) were used to avoid negative values in simulations. Functions for calculating separate ly for both data variables, $\ln(t)$ and $\ln(T)$, the means and standard deviations as well as correlations between these variables were needed. For simulation, a function generating random numbers from normal distribution was used and the simulated values for t and T could be calculated (see Paper VI for more details). To assess the probability of product being cooked within any t,T-segment the percentage of simulated data points (t,T) belonging to the specified t,T-segment was calculated.

An exponential dose-response model, which was also chosen for the recent FAO/WHO (2004) risk assessment study on RTE foods, was used in risk assessment:

$$P = 1 - e^{-rN},\tag{Eq. 3}$$

where P is the probability of an adverse effect, N is the number of biological agents consumed and r is a constant specific to each pathogen. The three r-values used in this study (presented in Buchanan et al., 1997; Linqvist and Westöö,

2000 and FAO/WHO, 2004) are presented in Table 1 of Paper VI along with the main characteristics of the models. The biological end point in all of these models was invasive listeriosis. The step-by-step presentation of calculating the risk with the exponential model using the *r*-value from the study of Buchanan et al. (1997) and worst-case point-estimates is presented in 4.7.3.

4. Results and discussion

4.1 Hygienically most problematic food processing equipment (Paper I)

The response rate for the questionnaire on equipment hygiene was 23.9% (44/184): 14 plants representing the meat, poultry and RTE food industry, 11 bakeries, 8 fish companies, 6 dairies and 5 plants from other branches of the food industry answered. The response rates obtained from both questionnaires (see also 4.2.1) are comparable with the rates of other recent postal surveys sent to the food industry (Hielm et al., 2006). There were most often 10–90 employees in the plants (69.2%) (n=39). The respondents (n=43) were most often working as quality managers or hygiene-responsible operatives (44.2%) or production managers (25.6%). As the equipment used by Finnish food producers is purchased mainly from other countries (mainly Europe), but also from Finland, the problems in hygienic design and the conclusions of this study are also applicable to other countries.

The respondents were asked to name the five hygienically most problematic pieces of equipment in their plant. There were 39 equipment choices given in the questionnaire and an option to mention other equipment as well. Altogether 61 types of equipment were mentioned at least once, which showed clearly that the hygiene problems in each company were specific. Packaging machines, conveyers, dispensers, slicing machines and cooling machines were considered the most problematic equipment (Table 11). In previous studies, packaging machines, conveyers, slicing machines and cooling machines have also been found to be a source of *Listeria* contamination (Humphrey and Worthington, 1990; Gravani, 1999; Miettinen et al., 1999a; Tompkin, 2002). The main reason was poor hygienic design. Self-made and domestic equipment were considered more hygienic than equipment purchased from outside Finland. This may be due to easier communication of problems and their solutions between manufacturers and food processors.

Packaging machines were cited most often as unhygienic equipment (Table 11). The shelf-life of a product depends to a great extent on proper packaging, the last step before the product is transported to retailers and to the consumers. A common problem with packaging machines is that they often do not tolerate water because of electronic circuits, which makes the cleaning and disinfection procedures very difficult.

Most pushlomatia	No. of responses						
equipment	All n=44	Meat and poultry <i>n</i> =14	Bakery n=11	Fish <i>n</i> =8	Dairy <i>n</i> =6		
Packaging machines	20	7	4	5	3		
Conveyors	17	7	6	1	3		
Dispensing equipment	15	4	8	2	1		
Slicing machines	15	9	3	3	_ ^a		
Cooling equipment	9	6	2	-	1		
Proving drawers	6	-	6	-	-		
Pasteurising equipment	5	-	1	-	4		
Cold stores	4	1	-	1	2		
Heat exchangers	4	1	-	-	3		
Pumps	4	1	-	-	2		
Filleting machines	3	-	-	3	-		

Table 11. The hygienically most problematic equipment in different food industry sectors (n = number of respondents).

^a "-" = not mentioned by the respondents

The respondents (n=30) stated that food processing equipment should be easy to dismantle and clean, and more simple constructions should be used to improve equipment hygiene. The materials should tolerate strong cleaning and disinfecting agents as well as heat. Furthermore, it should also be possible to use water in cleaning the equipment and process surfaces. Coverings should be easy to open for cleaning. These results stress the importance of including hygienic aspects in the equipment design at the beginning of the design process. However, the technical and occupational safety aspects must also be taken into account. If cleaning of the equipment is difficult due to these factors, the best possible solution for assuring cleanability, e.g. targeted cleaning programs to problematic sites, should be established.

Operational reliability (34%) was clearly the most important factor affecting the acquisition of equipment, followed by costs (20%) and cleanability (11%) (n=38). According to 43.2% of respondents, the manufacturers have provided instructions on how to clean the equipment and according to 40.9% some of the manufacturers have (n=44). When instructions were available, most respondents (76.7%) followed them (n=43), although 58.1% of respondents found the instructions inadequate. Many (38.1%) of the respondents (n=21) answered e.g. that equipment manufacturers are not interested in or do not understand the hygienic aspects of their equipment. However, manufacturers are required by the EU Machine Directive (89/392/EEC, revised 98/37/EC) to give instructions for cleaning the equipment. This Directive does not specify what kind of instructions should be included in the material. Many of the equipment manufacturers do not have sufficient knowledge of cleaning to give sufficient information to the client and because of this, instructions should be made in cooperation with cleaning specialists. There should also be cooperation on hygiene aspects between equipment and sanitizing agent manufacturers and food processors already at the design phase of the equipment.

4.2 Hygienic working practices of maintenance personnel (Paper I)

4.2.1 Significant aspects according to the questionnaire

In the questionnaire on hygienic working practices of maintenance personnel answers were obtained from 23.6% of the plants (n=106). Out of 778 employees 127 (16.3%) answered the mail survey. Of these, 59 were maintenance personnel and the rest (68) were food-handlers, quality managers and cleaning personnel. One third (33.9%) of the respondents worked in the meat and poultry industry, 26% in the bakery industry, 20.5% in the dairy industry, 1.6% in the fish industry and 18% in other sectors of the food industry.

According to this study, consumer complaints could rarely/seldom be linked directly to the work of maintenance personnel (62.5%, n=16). However, as pathogens such as *L. monocytogenes* can be transferred between processing surfaces and food products (Table 6), the hygienic practices of employees are of utmost importance in preventing contamination of equipment by these bacteria.

Most of the maintenance personnel worked in the production area continuously (61%) or at least visited the production area 5 times (23.7%) during a work shift and 42.4% reported having touched surfaces in contact with food often or always. Whereas they were conscious of this and they knew (91.5%, n=59) which surfaces came into contact with food, only 55.9% of them usually wore gloves when working in the food production area and even fewer washed their hands in situations where they should in order to work hygienically (13.6% after smoking and 23.7% before starting to work without gloves). One reason for not wearing gloves is probably that they hinder the performance of some work tasks. In situations like this the importance of washing of hands should be highlighted.

Almost all (89.8%) of the maintenance personnel had personal tools which they themselves cleaned (n=59). One third (32.2%) of the respondents answered that they washed their tools once a day or always after work. The others answered that they washed their tools more seldom and 32.2% only once a year or never. The majority of maintenance personnel and quality managers (69%, n=87) considered that there were enough washing points and adequate cleaning agents and disinfectants available (74.7%, n=83). For tools that are in common use, no specified persons were in charge of the cleaning in 71.9% of all the cases reported (n=57).

According to the majority (88.2%, n=59) of maintenance personnel, foreign bodies were never or seldom left on surfaces after the maintenance work but, according to the majority of food and cleaning personnel (68.2%, n=66), they were sometimes or even often left. Clear differences between opinions were identified. The responsibility for cleaning the equipment after work by the maintenance personnel was not always defined. All responsibilities in hygiene matters in food companies should be clear.

According to the maintenance personnel and quality managers, most of the maintenance personnel had written hygiene rules given by the food company (64.2%) or by the maintenance company (32.3%) (n=81). Most (63.6%) of the maintenance personnel knew where to find the hygiene rules (n=55). It is very important that the rules are available for all. These rules must be clearly written and should also include the maintenance personnel and their work.

Meetings between maintenance, food-processing and cleaning personnel in which hygiene issues were discussed, were not arranged in 52.9% of the plants (n=119). One third (33.9%) of maintenance personnel reported that they had not received sufficient information about hygiene aspects commensurate with their work. Studies on food hygiene were not included in their basic education (89.9%, n=59). The requirement for sufficient hygiene knowledge of especially this personnel group should also be stated in legislation and studies on food hygiene should be obligatory in their basic education. As publications on hygienic working practices of maintenance personnel have not previously been available, the results can only be compared with results of studies made on hygienic practices of food employees. Several deficiencies were also detected in these studies (Table 5).

4.2.2 Microbiological sampling in food processing

The occurrence of *L. monocytogenes* and the total number of aerobic bacteria in samples taken from the tools, environment and personnel (protective clothing and hands of maintenance personnel) are presented in Table 12. *Listeria* spp. was found in six samples (8.5%), of which *L. monocytogenes* was found in only one sample (1.4%) taken from a screwdriver. Tools contained clearly (p<0.05) less aerobic bacteria than samples from personnel and environment. Samples from the personnel contained on average more bacteria than samples from the environment, but the difference was not significant (p>0.05). *Enterobacteriaceae* were only found on one of the gloves (100 CFU/cm²).

Although *L. monocytogenes* was not found from the samples taken from clothes, the amount of aerobic bacteria on maintenance personnel clothes and tools was on average high (Table 12). This result pinpoints the need for regular and sufficiently frequent changing of the protective clothing and the need to change them whenever they get dirty. As *L. monocytogenes* was sampled from one screwdriver and another *Listeria* spp. from a pneumatic machine in common use (see Paper I) tools are potential vectors of *L. monocytogenes* transfer within food processing. This emphasizes the need to clean and disinfect them regularly, preferably daily and always whenever they get soiled.

Table 12. Results of microbiological samples taken from clothes, tools and work environment of the maintenance personnel. The samples were taken with gauze pads for Listeria spp. and with PetrifilmTM plates for aerobic bacteria sampling.

	Listeria spp.	ria spp. Aerobic total bacteria					
Samples	No. of positive samples / No. of samples	No. of samples	Range (CFU/cm ²)	Average (CFU/cm ²)	St. dev. (CFU/cm ²)		
Tools	2/40 ^a	50	1–40	5	8		
Personnel	4/24	36	1-100	14	23		
Environment	0/7	9	3–20	8	5		

^a L. monocytogenes was detected from one sample taken from a screwdriver.

4.3 Discriminatory power of automated ribotyping compared with PFGE in distinguishing *L. monocytogenes* isolates (Paper II)

Identification of L. monocytogenes isolates to strain level by typing is needed in food plants for tracing contamination routes and identifying possible plantspecific strains causing problems. Rapid tracing of L. monocytogenes contamination sources is of utmost importance in the prevention of product contamination. Hitherto the only fully automated typing method is automated ribotyping and thus the discriminatory power of this method was compared to the generally recognized method PFGE and the traditionally used serotyping. Digestion with *Eco*RI generated 16 different ribogroups or ribotypes (RTs) from the 486 L. monocytogenes isolates and when 121 isolates representing all the 16 RTs were further typed with PvuII, 19 RTs were generated (Table 2, Paper II). When PFGE was used, digestion with AscI resulted in 42 and with SmaI in 24 macrorestriction patterns from the 121 isolates analyzed. When the patterns were combined, 46 final PFGE types were generated. The *DI* was 0.878 for ribotyping with EcoRI and 0.867 with PvuII. The overall DI for ribotyping with both enzymes was 0.906. In the case of PFGE, the DI of typing with AscI was 0.960 and with Smal 0.920, the overall DI being 0.966. Most of the isolates (109/121) belonged to the serotype 1/2 (Table 2, Paper II). Only four of the isolates were of serotype 4b (3.3%) of the isolates) and two of serotype 3 (1.7%). Some isolates (6/121) were not typeable with the commercial kit used.

From the isolates of four plants (E[29 isolates], I[7 isolates], J[25 isolates], and P[25 isolates]), PFGE generated more PFGE types (8, 5, 11 and 14, respectively) than ribotyping RTs (5, 4, 7 and 9, respectively). PFGE could further divide 10 RTs into a total of 33 PFGE types, indicating more genetic types from different plants than ribotyping. By contrast, ribotyping could also divide 3 PFGE types into a total of 6 RTs, indicating higher discrimination between these isolates than that obtained by PFGE.

Thus the discrimination power is not only dependent on the method, but also on the isolates and enzymes used. With automated ribotyping it is possible to ribotype the first pure culture colonies without any conventional time-consuming preliminary tests in 8 h. More expertise and careful manual work is needed for running PFGE. With PFGE, identification of an isolate from pure culture can be performed in 30 h (Graves and Swaminathan, 2001).

According to the results presented above (see Paper II), as well as in Paper VI, PFGE had higher discriminatory ability than automated ribotyping. However e.g. in the Paper VI, the strain occurring frequently during one year in products of one producer, and surviving milder heat treatment, could be distinguished from the other isolates with both methods. Automated ribotyping has successfully been used for investigating contamination sources in food plants (Suihko et al., 2002), and mainly based on its automation and rapidity, it is a good tool for screening large numbers of isolates in contamination studies. However, in cases of outbreaks, the identical patterns must be confirmed by PFGE.

4.4 Survival and transfer of *L. monocytogenes* in lubricants

4.4.1 Hygiene of lubricants according to the questionnaire (Paper I)

Lubricants are needed for various purposes in almost all equipment to maintain proper functioning (see 1.1.3). The questionnaire on equipment hygiene (3.1) also contained questions on hygiene of lubricants used in food plants. 21.4% of the respondents (n=42) had noticed hygiene problems in the use of lubricants. The respondents (n=11) gave the following reasons for hygiene problems in lubricants: lubricants collect a lot of soil (72.7%); traces of lubricants are left on

production surfaces after maintenance work (63.6%); inability to clean surfaces from lubricant residues (54.5%); and high levels of micro-organisms being found in the samples taken from sites containing lubricants (36.4%). A few earlier studies also showed the ability of lubricants to support the growth of micro-organisms (van der Waa, 1995; Rossmoore, 1988; Ortiz et al., 1990; Hamilton, 1991).

4.4.2 Survival of L. monocytogenes in lubricants (Paper III)

Only a few investigations have been available on the occurrence and survival of any micro-organisms, not to mention *L. monocytogenes*, in lubricants used in the food industry. The ability of *L. monocytogenes* to survive in lubricants used in food processing equipment was investigated by inoculating bacteria to lubricants and taking samples during a 14 d test period. The amount of *L. monocytogenes* in lubricants decreased significantly (p<0.05) during the 14 d test period except in the case of synthetic conveyer-belt lubricant B and rapeseed oil D, in which the amount even marginally (p>0.05) increased at 20 °C. Because of the clear survival of *L. monocytogenes* in rapeseed oil, it would not be recommendable for lubricating purposes in the food industry unless it were to be cleaned off the surfaces and replaced with new oil daily. *L. monocytogenes* also survived well in the synthetic conveyer-belt lubricant B diluted in water, which clearly indicates that water enhanced the survival of *L. monocytogenes*. The dry synthetic lubricant proved to be a better choice for lubricating conveyer belts.

Clear differences in survival of *L. monocytogenes* in different lubricants were observed. The following pure lubricants reduced the amount of *L. monocytogenes* >3 log CFU/g during the 14 d incubation period at room temperature: mineral-oil based hydraulic oil E, dry synthetic conveyer-belt lubricant F, mineral-oil based multipurpose grease I and chain lubricant C (Fig. 1a, Paper III). In these lubricants the reduction was statistically significant (p<0.05) already after 24 h. Two lubricants (the white-oil based aluminium complex grease J [USDA H1] and synthetic multipurpose grease K) killed *L. monocytogenes* in both pure and soiled conditions both at room (20 °C) and refrigerated (5 °C) temperatures already 0.5 h after inoculation. Thus, the use of these lubricants can be recommended.

Cold temperature and soiling had different effects on each lubricant. The listericidal effect of the dry synthetic conveyer-belt lubricant F and mineral-oil based hydraulic oil E (both pure and soiled) was reduced (p<0.05) when the temperature was decreased from 20 °C to 5 °C. Conversely, low temperature clearly (p<0.05) increased the listericidal effect of chain lubricant C (Fig. 1b, Paper III). Soiling reduced the listericidal effect of chain lubricant C used as chain lubricant and mineral-oil based hydraulic oil E especially at 5 °C, but it did not have a clear effect (p>0.05) on the lubricants in general (not shown in Figs.). Organic material has been reported to reduce the listericidal effects of sanitizers and lubricants reduced the amount of *L. monocytogenes* better (p<0.05) than food-grade lubricants, but use of food-grade lubricants is required in food contact areas.

Although the amount of *L. monocytogenes* in most lubricants, both pure and soiled, decreased significantly (p<0.05) during the 14 d test period, lubricants may act as a source of contamination on the basis of the results obtained on the survival of *L. monocytogenes*. Water in lubricants should be avoided. In food manufacturing processes the change period for lubricants is often longer than two weeks, which may result in higher quantities of soil and better survival and growth of *L. monocytogenes* than detected in this study. Suitability for use in different temperatures and in clean or soiled conditions should also be considered from the microbicidal point of view.

4.4.3 Transfer of *L. monocytogenes* from lubricants to stainless steel surfaces and vice versa (Paper III)

The ability of lubricants to act as vectors for transferring *L. monocytogenes* was investigated by studying the transfer of *L. monocytogenes* from lubricants to stainless steel surfaces and vice versa. Significant (p<0.05) transfer of *L. monocytogenes* from lubricants to the surfaces of stainless steel discs was detected (Table 3, Paper III). The largest amount of *L. monocytogenes* (CFU/g) was detected on surfaces of discs which were incubated in soiled grease lubricants A and G. *L. monocytogenes* bacteria were also well transferred to the stainless steel surfaces from synthetic conveyer-belt lubricant B and the rapeseed oil D.

Significant (p<0.05) transfer of *L. monocytogenes* from stainless steel surfaces to lubricants at room and refrigerated temperature was seen in the case of both conveyer-belt lubricants, B and F (Table 4, Paper III). No transfer into synthetic grease lubricant A containing Teflon (USDA H1) was detected during the 24 h follow-up study. *L. monocytogenes* was transferred >4 log CFU/g to mineral-oil based hydraulic oil E at 5 °C (strain III) (Table 4, Paper III) and sporadic transfer to other lubricants (C, D and G) was detected. No previous studies on the transfer of bacteria from surfaces to lubricants or vice versa have been reported. The results indicate that lubricants, especially conveyer belt lubricants, may act as contamination vectors between processing surfaces. The lubricants should be regularly changed and the surfaces cleaned and disinfected (see 4.5) before adding new lubricant. The requisite changing frequency should be determined by monitoring the microbiological contamination level by sampling.

4.5 Susceptibility of *L. monocytogenes* to disinfectants (Paper IV)

4.5.1 Efficacy of disinfectants in suspension

Successful daily cleaning and disinfection are needed as means of decontamination of food processing equipment and premises for producing safe foods. As sanitizing is usually performed in cold premises, the efficiencies of commonly used disinfectants were investigated at +5 °C. All the disinfectants (Table 10), except alkaline hypochlorite-containing disinfectant E, were effective (ME > 5) in both clean and soiled conditions in suspensions. However, the peracetic acid-based disinfectant A in both clean and soiled suspensions, as well as the peracetic acid-based disinfectants C and H in soiled conditions failed to inactivate some strains at the lowest recommended concentrations given by the manufacturer (Table 3, Paper IV). When the concentrations were increased and the duration was extended to 10 min, the three latter disinfectants were also effective.

Tuncan (1993) stated that cold temperature did not affect the efficacy of chlorine. However, increased effectiveness due to an increase in temperature has also been reported (El-Kest and Marth, 1988b; Orth and Mrozek, 1990). The alkaline hypochlorite-containing disinfectant E was also tested as recommended

by the manufacturer at 20 °C. However, the increase in temperature and an increase of the concentration from the lowest recommended (0.05%) in-use concentration to 0.2% using a duration time of 10 min was not sufficient to inactivate all the strains. The pH of in-use concentrations of agent E was between 8 and 10. According to McDonnell and Russell (1999) in aqueous solution between pH 4 and 6, chlorine exists predominantly as hypochlorous acid (HCLO), the active moiety, whereas above pH 9, the hypochlorite ion (OCl-) predominates. The pH of this disinfectant should be lower to increase its microbicidal efficiency.

4.5.2 Efficacy of disinfectants on surfaces

As attachment of the cells to surfaces has previously been shown to reduce the efficacy of disinfectants (see 1.4.2), the efficacy of disinfectants was tested also on surfaces. In this thesis, all the disinfectants (Table 10) were effective (ME>4) when applied to stainless steel and PE surfaces, except the QAC-based disinfectant F and the peracetic acid-based disinfectant H against one strain (Table 5, Paper IV). The QAC-based disinfectant F was ineffective (ME<4) against two of the strains on both surfaces. Previously cold temperature has also been shown to reduce the efficacy of QACs and of iodofor on *Listeria* spp., especially at low concentrations (Orth and Mrozek, 1990; Tuncan, 1993). However, the recommended duration for this QAC-based product to act was 0.5–1 h. The alkaline hypochlorite-containing disinfectant E was effective on the surface, which may be explained by the fact that *L. monocytogenes* was growing on the surfaces on the agar plate covered with a filter paper and the amount of protein on the surface was low.

According to Taormina and Beuchat (2001), heat resistance of *L. monocytogenes* was increased after exposure to alkali. The safety of foods requiring heat treatment may be endangered if they are contaminated by *L. monocytogenes* cells which have survived exposure to processing environments ineffectively cleaned or sanitized with alkaline detergents or disinfectants. Therefore the efficiency of cleaning and sanitizing treatments should be assured at the plant level, e.g. in cooperation with research laboratories, in the conditions prevailing in the process, and residues of the agents should be rinsed off the surfaces completely.
4.6 Transfer of *L. monocytogenes* during slicing of coldsalted salmon (Paper V)

As recontamination has been identified as a significant cause of food contamination, information on the level of recontamination for risk assessment purposes is needed. Slicing machines are one of the hygienically most problematic pieces of equipment in the food industry (Table 11). As cold-salted salmon is a RTE product, which has been associated with contamination by *L. monocytogenes* in several studies (Table 5), potential cross-contamination from a contaminated blade to uncontaminated cold-salted salmon slices was simulated.

4.6.1 Transfer of *L. monocytogenes* from slicing blade to slices

Transfer of L. monocytogenes was calculated over a total of 39 slices. There was a progressive exponential reduction in the quantity of L. monocytogenes transferred (Fig. 1). When compared to the inoculum level of the blade, clearly (p < 0.05) lower total numbers of L. monocytogenes were transferred when the inoculum level was lower, the temperature was colder or the attachment time was longer compared to the experiment made at room temperature with a high $(8.4\pm0.4 \log \text{ CFU/g})$ inoculum level and a short attachment time (10 min). For example $5.3\pm0.3 \log \text{CFU/g}$ was transferred to the second slice when the inoculum level was 8.4±0.4 log CFU/blade and the amount was reduced ca. 1.6 log CFU/g during slicing of 39 slices (Fig. 2a, Paper V). Based on the result data of the samples (2-39), when compared to the inoculum level of the blade, there were no statistically significant differences in the logarithmic reduction of L. monocytogenes numbers in slices between the different experiments (p>0.05), although a marginally lower reduction was detected at 0 °C compared to room temperature. The transfer percentage (the amount of L. monocytogenes in all 39 slices compared to the inoculum) varied between 0.00011 and 0.17%, being lowest at 0 °C. However, when calculated with the predicted values, instead of using the results of 39 slices directly, a significantly (p < 0.05) lower logarithmic reduction in the number of L. monocytogenes between slices was found when slicing was at 0 °C compared to the experiment made at room temperature with a high inoculum level and a short attachment time, indicating that cold temperature prolonged the transfer. However, in all experimental conditions, the number of bacteria decreased quite rapidly (i.e. after slicing the fourth fillet) to <1 log CFU/g.

The reduction in quantity of L. monocytogenes transferred was lower than that reported by Vorst et al. (2006a) for turkey breast, bologna and salami (2 log CFU/20 slices). Soft salmon material (salmon fillets consisting mainly of protein, fat and moisture) on slicer surfaces as well as solidification of fat most probably slowed the transfer at colder temperatures. According to Midelet and Carpentier (2002), who studied transfer of bacteria including L. monocytogenes from various materials to pieces of beef, in most cases the inoculation concentration had the strongest influence on the total number of CFU detached. The attachment strength of bacteria on different materials also had a significant (p < 0.05) effect on transfer. In the current study, based on the predicted values higher numbers (p < 0.05) of L. monocytogenes were also transferred to slices from the slicing blade when the inoculum was higher or when the attachment time was shorter. When the blade inoculum was low, ca. 3 log CFU/blade, only a few colonies were found on the first 10 slices and no colonies were detected on the slicer surfaces, including the blade. Levels below the agar detection limit were most probably present but were not examined by enrichment.

When a salmon fillet was inoculated with *L. monocytogenes* (surface inoculum of fillet was 7.6±0.1 log CFU *L. monocytogenes* per fillet) and sliced, the blade contained 3.9 ± 0.8 log CFU/10 cm² (i.e. 5.5 ± 2.4 log CFU/blade). 3.2 ± 0.4 log CFU/g was initially transferred to the subsequent sliced uninoculated fillet and the overall reduction in transfer was 1.5 log CFU after 39 slices. The reduction was only marginally (*p*>0.05) lower compared to reduction when the blade was initially contaminated. When the predicted values were used with higher (156) slice numbers (Fig. 1), significantly smaller logarithmic reduction was observed compared to the experiment, in which the blade was directly inoculated (8.4±0.4 log CFU/blade, 10 min) at room temperature.

The contamination level on the holding plate and cover depended on the level of *L. monocytogenes* on slices, especially in experiments made at room temperature, i.e. the higher the level was on slices, the higher the level was on equipment surfaces (Table 1, Paper V).

The roughness (A'/A) of the slicing blade was 2.3 ± 0.1 on the sharp side and 1.7 ± 0.1 on the flat side in the beginning of the experiments, and 2.0 ± 0.2 and 2.7 ± 0.6 , respectively, at the end of the experiments. The results and pictures taken of the new blade and the blade at the end of the experiments (pictures not

shown) by reflection confocal microscopy clearly demonstrated blade wear over the course of the experiments, emphasizing the importance of regular maintenance of the blade in food processing operations. The blade, the blade guard and holding plate should be periodically dismantled, cleaned and sanitized to avoid attachment of and prolonged product contamination with *L. monocytogenes*. Additionally, the blade should be sharpened regularly.

4.6.2 Model of transfer of *L. monocytogenes*

Based on the results of the transfer experiments performed in the laboratory, transfer in different conditions was modelled. An exponential model of transfer as a function of slice number provided a reasonable fit across all treatments, e.g. average $r^2>0.7$, except that in the 0 °C study the r^2 -value was 0.63:

$$y=a^*e^{(-x/b)}; (Eq. 4)$$

The values for constants a and b are different for each experiment. The model describes a microbiological decay curve as a function of slice number. The results of the predictions are most reliable when the data is obtained from experiments made in conditions prevailing in the process where the model will be applied (e.g. temperature, attachment time of inoculum, type of slicer etc.).

In this thesis, limited data from microbiological analysis was used to assess transfer of *L. monocytogenes* among processing surfaces and product. According to the results, *L. monocytogenes* was readily transferred from contaminated slicing blade to cold-salted salmon slices. The highest numbers of *L. monocytogenes* on slices were detected in the beginning of the slicing (except in the very first slice). Based on the results, the expected numbers of *L. monocytogenes* on slices due to recontamination from the blade are low, especially after slicing the first slices. The results obtained in this study can be used to assess the level of recontamination in the salmon process and they also provide an example how to estimate the impact of recontamination on other products.



Figure 1. Predicted transfer of L. monocytogenes from inoculated blade to uninoculated salmon fillets during slicing of 200 slices in different conditions. After slice 39 the predictions are extrapolations.

4.7 Producer level quantitative risk assessment of *L. monocytogenes* (Paper VI)

4.7.1 Prevalence and numbers of *L. monocytogenes* in broiler legs

Information of prevalence and levels of *L. monocytogenes* in raw marinated broiler legs was collected for performing a producer level risk assessment (4.7.3). Overall, 39% (72/186) of the samples were contaminated with *L. monocytogenes* (Table 13) (36% of the samples stored at 6 °C and 57% of the samples stored at 10 °C). The prevalence of *L. monocytogenes* varied between the producers, being 18% (11/62) for producer A, 69% (43/62) for B and 29% (18/62) for C (see Paper VI). The numbers of *L. monocytogenes* were low, between 0.3 and 147 CFU/g in all the samples stored at 6 °C and between 0.5 and 300 CFU/g in all the samples stored at 10 °C.

In Finland prevalences between 35% and 62% have been reported for retail raw broiler meat (Miettinen et al., 2001; Johansson et al., 2003). In other countries levels of *L. monocytogenes* positive samples in raw broiler meat have been similar (Cox and Bailey, 1999). Only a few investigations are available on the

levels of *L. monocytogenes* in raw broiler or chicken meat. According to Mead et al. (1990) and Rørvik and Yndestad (1991), raw chicken normally contains *L. monocytogenes* $<10^3$ CFU/g. Our results are thus in agreement with these previous findings. Franco et al. (1995) reported levels of $>10^3$ CFU/g for raw chicken leg skin and leg muscle meat in many samples taken directly from a poultry processing plant, and found them most contaminated from the poultry pieces.

Table	e 13.	Prev	alenc	e and	levels	of L.	mono	cytog	genes	s (L.1	n.) <i>i</i>	n m	arina	ted	broil	er
legs	obta	ined j	from	retail	shops	and	stored	at b	б°C	or a	t 10	°C	until	the	end	of
their	shelj	f life.														

Producer	Production period	No. of <i>Listeria</i> spp. positive samples (%)	No. of <i>L. m.</i> positive samples (%)	No. (CFU/g) of <i>L. m.</i> Range (Average)
А	Feb/Mar – Nov/Dec -02	26/55 (47)	9/55 (16)	0.3–29.7 (5.1)
	Jan/Feb ^a -03	2/7 (29)	2/7 (29)	0.5–5.0 (2.8)
В	Feb/Mar – Nov/Dec -02	40/55 (73)	37/55 (67)	0.3–147 (7.4)
	Jan/Feb ^a -03	6/7 (86)	6/7 (86)	2.0–180 (39.2)
С	Feb/Mar – Nov/Dec -02	14/55 (25)	14/55 (25)	0.3–2.7 (0.9)
	Jan ^a -03	4/7 (57)	4/7 (57)	0.5–300 (84.6)
	Total	92/186 (49)	72/186 (39)	0.3–300 (12.7)

^a stored at 10 °C

4.7.2 Thermal inactivation of *L. monocytogenes* by heating

The temperatures used in the laboratory experiments were based on the consumer tests. For cooking the broiler legs, the consumers used 43–105 min (average 64 min) and the oven was set to 175 °C, 200 °C, 205 °C or 225 °C. The lowest average temperature detected in consumer ovens during cooking was 132 °C. 74% (14/19) of the consumers kept the broiler legs longer than 50 min in the oven, which was the producer's recommendation for cooking at 175 °C.

The consumer cooking data was divided into four (t,T) segments (A–D) with (A) \geq 50 min, \geq 175 °C; (B) \geq 50 min, between \geq 132 °C and <175 °C; (C) \geq 50 min, <132 °C; (D) <50 min, any temperature (Fig. 2, Paper VI). When calculated using a simulation as presented in 3.6.5, the corresponding simulated probabilities were: *P*(A) = 0.25, *P*(B) = 0.51, *P*(C) = 0.03 and *P*(D) = 0.21. However, we limited our example to calculating conditional risk when consumers heat the broiler legs according to producer's instructions, for \geq 50 min. Accordingly a simulation was performed for this data to obtain the conditional probabilities *P*(*t*,*T* | *t*>50 min). Thus,

P(T > 175 | t > 50 min) = 0.25 / (1 - 0.21) = 0.32 $P(132 < T \le 175 | t > 50 \text{ min}) = 0.51 / (1 - 0.21) = 0.64$ $P(T \le 132 | t > 50 \text{ min}) = 0.03 / (1 - 0.21) = 0.04.$

In laboratory experiments, of the 21 samples in each of the two experiments (132 °C 50 min and 175 °C 50 min), 12 (57%) were contaminated with *L. monocytogenes* from <1 to 300 CFU/g. The heating eliminated *L. monocytogenes* from all the samples, except for one sample at 132 °C. The level of *L. monocytogenes* in this sample before heating was 22 CFU/g. The worst-case cooking scenario was 132 °C \geq 50 min, as none of the consumers in this study actually used the combination of \leq 132 °C \leq 50 min (see Fig. 2, Paper VI).

4.7.3 Quantitative risk assessment

Quantitative risk assessment was performed on the basis of the contamination (4.7.1) and heating study (4.7.2) data (see Fig. 3, Paper VI). The worst-case probability of a sample being *L. monocytogenes* -positive after heating would be calculated using the probabilities calculated in 4.7.2 as follows:

 $P(L. m.-\text{positive after heating} | L. m.-\text{positive before heating and } t > 50) = 0.32 \times 0 + 0.64 \times (1/12) + 0.04 \times 1 = 0.09.$

As the worst-case probability, the producer could choose the highest % of positive samples obtained during the time period of interest. In our study we used 39%, the value obtained from the samples, as the worst-case value. Then,

P(L. m.-positive after heating; worst) = P(L. m.-positive samples)before heating; worst) × P(L. m.-positive after heating | L. m. positive before heating and t > 50 = 0.39 × 0.09 = 0.04. (Eq. 5)

L. monocytogenes was only detected after cooking but, because the number of *L. monocytogenes* in broiler leg in which the bacteria survived was 22 CFU/g before cooking, this number was used as the worst-case number of *L. monocytogenes* after cooking. When this number was multiplied by the greatest weight of a broiler leg (456 g), we obtained a worst-case approximation of the level of exposure/portion/individual, $N(\max)$, of 1.032×10^4 CFU, with the assumption that one portion is one broiler leg. According to Eq. (3), the worst-case dose-response from one broiler leg would then be:

 $P(\text{listeriosis from } L. m.\text{-positive broiler leg; worst}) = 1 - e^{(-1.18 \times 10^{10} \times 1.032 \times 10^4)} = 1.18 \times 10^{-6}.$

As 6.3 million kg of marinated broiler legs were consumed in Finland, the number of consumed legs was 6.3 million kg / 0.301 kg / 5.3 million consumers (population of Finland) = 3.95 marinated broiler legs/person/year. Of the population of Finland, 16-22% belong to risk groups (Maijala et al., 2001). As a worst-case estimate, 22% of the population was estimated to belong to the risk group. Thus the worst-case listeriosis risk from heated marinated broiler legs for people belonging to a risk groups would be:

 $P(\text{listeriosis; risk group, worst}) = P(L. m.\text{-positive after heating;}) \times P(\text{listeriosis from } L. m.\text{-positive broiler leg; worst}) \times P(\text{risk group; worst}) = 0.04 \times 1.18 \times 10^{-6} \times 0.22 = 9.14 \times 10^{-9}. \quad (\text{Eq. 6})$

When the number of portions eaten/person/year is included, this gives the expected cases of listeriosis as $9.14 \times 10^{-9} \times 3.95$ portions/person/year $\times 5.3$ million = 0.19 cases/year.

The risk for the general population was also calculated (see Table 14) as well as values obtained by using the following average point estimates: level of *L. monocytogenes* -positive broiler legs 34% (estimated mean level, see Paper VI); P(L. m.-positive after heating | *L. m.* -positive before heating and t>50) = $(0.32 \times 0 + 0.64 \times (1/12) + 0.04 \times 1)/2 = 0.05$, which assumption was made

based on the use of a time × temperature combination of 50 min × 175 °C or more by 15/19 of the consumers; number of *L. monocytogenes* after heat treatment, 2 CFU/g (estimated average concentration in raw marinated broiler legs see Paper VI); average weight of a broiler leg portion, 301 g; and the portion of the total population belonging to the risk group (16 + 22)/2%=19%. The results obtained with different *r*-values of dose-response models are presented in Table 14.

Table 14. Expected cases of listeriosis/year from heated marinated broiler legs when heated according to the producer's instructions, based on different dose-response models.

	No. of listeriosis cases/year							
Dose-response model used	High-risk j	population	General population					
	average	worst-case	average	worst-case				
Buchanan et al., 1997	4.32×10^{-3}	0.19	_a	-				
Linqvist and Westöö, 2000	2.05×10^{-2}	0.91	-	-				
FAO/WHO, 2004	-	-	8.34×10^{-6}	3.07×10^{-4}				

^a "-" = No dose-response model available.

According to the results of the point estimate-based risk assessment $4.32 \times 10^{-3} - 2.05 \times 10^{-2}$ (on average) people belonging to the risk group in Finland would annually be at risk of suffering severe listeriosis from marinated broiler legs after cooking according to the instructions given by producers. As the risk was negligible for the general population, the results of the point estimate-based risk assessment indicated that the risk is very low. At the plant level it should be considered which products require a thorough risk assessment. In addition, factors causing uncertainty in risk assessment must be described in order to be able to evaluate the level of confidence and to avoid unrealistic scenarios (see description in Paper VI).

The producer can affect the prevalence and numbers of *L. monocytogenes* on a product before it leaves the factory, and adjust the sell-by date and the guidelines for cooking it. Using this kind of risk-based approach, the effects of different management actions on the number of cases of listeriosis can be estimated and thus a more comprehensive understanding of the risk to consumers obtained. A similar approach can be used to assess the risk of *L. monocytogenes* and other microbiological hazards for different products.

Additionally, the results indicate that Finnish oven-cooked marinated broiler legs do not present a significant risk from *L. monocytogenes* if recontamination after heating is avoided. Recontamination can be a major issue in contamination of the processed products (see 1.3.6) and should be prevented. This emphasizes once again the importance of good equipment hygiene.

5. Conclusions

Occurrence of *L. monocytogenes* in different food products is a universal problem in the food industry as the bacterium is widely distributed and tolerant against various environmental conditions. *L. monocytogenes* is known for its ability to persist in food processing equipment and therefore hygiene should be optimized against this pathogen. In this thesis, deficiencies in current process hygiene measures and improvements for minimizing the occurrence of *L. monocytogenes* were investigated and the following conclusions were drawn:

1. Equipment design and maintenance practices in food processing

Packaging machines, conveyers, dispensers, slicing machines and cooling machines were found to be the hygienically most problematic equipment in food processing plants. The main reason was generally poor hygienic design. In previous studies, these machines have also been found to be sources of *L. monocytogenes* -contamination. Thus equipment designers should focus their performance on more suitable equipment design. Additionally, their training in this topic must be increased.

The effect of hygienic practices of maintenance personnel on equipment hygiene and on occurrence of *L. monocytogenes* in processing equipment has often been a topic in food safety discussions but has not been studied to date. In the current thesis it was found that the maintenance personnel in food processing plants is a potential source of contamination of food processing equipment, and thus food products. Clear deficiencies in hygiene performance of this employee group were found. An indication was also found that *L. monocytogenes* could be transferred through maintenance work. Training of maintenance personnel in hygiene matters must therefore be increased and included in the legislation.

2. Automated ribotyping and PFGE in identification and tracing of *L. monocytogenes* in food processing

Typing of *L. monocytogenes* -isolates is needed in tracing contamination sources in food plants. The results of this thesis showed that PFGE had a higher discriminatory power for *L. monocytogenes* isolates than ribotyping. However, due to its automation and rapidity, automated ribotyping can be considered a good method for *L. monocytogenes* control purposes and for detecting contamination sources in food processing. In epidemiological studies, the results obtained with the automated ribotyping system must be confirmed by PFGE.

3. Lubricants and cold disinfectant treatments as vehicles in *L. monocytogenes* contamination routes in the food industry

Recently an ISO-standard and guidelines on Good Manufacturing Practices for manufacturing and use of food-grade lubricants have been published. However, the microbiological quality of lubricants has received little attention. The studies made in this thesis showed that *L. monocytogenes* survived during a 14 d experiment period in 9 out of 11 lubricants used in food-processing equipment. Clear differences between lubricants were found. It was shown, for the first time, that *L. monocytogenes* was transferred to stainless steel surfaces from lubricants and in many cases from surfaces to lubricants, indicating that lubricants may act as vectors in *L. monocytogenes* contamination from one surface to another. In addition to the requirements of toxicological safety and good technical performance, based on this study, prevention of survival and growth of bacteria should be considered when choosing lubricants for maintenance of food-processing equipment. Moreover, samples for detection of *L. monocytogenes* should be regularly collected from lubrication points of food processing equipment.

The eight commercial disinfectants were in general efficient against *L. monocytogenes* strains at +5 °C at the concentrations and effect times recommended by the manufacturer. Thus they were suitable for control of *L. monocytogenes* at the plant level, with only a few exceptions. The results pinpoint the need to use appropriate concentrations and duration, especially

in the case of QAC and hypochlorite-based agents. In some cases the lowest recommendations given by the manufacturer may not be sufficient. The efficiency of cleaning and sanitizing treatments at production plant level should be confirmed, e.g. in cooperation with research laboratories, under conditions prevailing in the processes.

4. Risk assessment at the food processing plant level

Information on the level of recontamination of products is needed for risk assessment purposes. An example of how predictive modelling and limited data from microbiological analysis can be used to assess the level of recontamination was given in the thesis. To assess recontamination, transfer of *L. monocytogenes* from slicing blade to slices of cold-salted salmon was investigated and clearly observed. An exponential model was suitable for predicting the expected number of *L. monocytogenes* (log CFU/g) on the salmon slices. In all test conditions, the numbers of *L. monocytogenes* were predicted to be reduced to <1 log CFU/g after the fourth uninoculated fillet was sliced. Salmon processors can use the results as a guide in risk management decisions.

A robust quantitative risk assessment of *L. monocytogenes* for one product was presented. Marinated broiler legs were used as an example product. This approach helps food processors in illustrating the risks caused by the products for consumers by estimating the effects of different risk management actions on the number of cases of listeriosis. This information can be utilized when planning risk management actions for own-checking systems. The results indicate that Finnish oven-cooked marinated broiler legs are not a significant *L. monocytogenes* risk, if recontamination after cooking is avoided. The results of the studies presented in this thesis also emphasize that risk assessment may be needed especially in case of products where recontamination after processing may typically occur.

Efficient control of *L. monocytogenes* at plant level is a combination of good equipment hygiene including functioning Good Hygiene Principles (GHP's) used by all employee groups, and rapid methodology in detecting contamination sources, as well as efficient hazard analysis systems utilizing

a risk assessment approach. Optimally hygienic solutions of the food plants for control of *L. monocytogenes* and other pathogens should be included in the design phase of the food processes and processing equipment as early as possible. They should be optimized for the conditions prevailing at each processing plant, which emphasizes cooperation between food processors, research institutes and suppliers of hygiene solutions. The measures should be environment- and energy-saving, which requires a high standard of equipment design. New design solutions, and materials as easy to clean and disinfect as possible, should be developed for reduced need of sanitizing and maintenance, e.g. changing parts of machinery. As rapid tracing of contaminants is a key issue in prevention of food-borne illnesses caused by contaminated products, development of rapid, reliable and easy-to-use automated methods for detecting contamination sources in food processes is needed.

The impact of different maintenance procedures on transfer of *L. monocytogenes* inside food plants should be studied further. Studies on the importance and mechanisms of recontamination have been launched during recent years. Several factors regarding transfer of *L. monocytogenes* need further investigations, including the effect of strain variability and product composition. Additionally, transfer from production scale slicers should be studied. Investigations on bacterial levels in lubricants used in maintenance of food processing equipment and their role in contamination of the equipment should be performed. Moreover, it should be investigated whether lubricants lead to the development of resistance and cross-resistance of *L. monocytogenes*.

Future risk assessment at the plant level should have a more comprehensive, quantitative approach with the assistance of computer programs. For this purpose, there is a clear need for additional quantitative data on *L. monocytogenes* contamination levels in different foods and production equipment.

References

Aalto, T., Häkkinen, M., Hallikainen, A., Hatakka, M., Johansson, T., Kostamo, P., Kuronen, H., Lahti, E., Lumme, M., Nuppunen, M., Pirhonen, T. and Törmä-Oksanen, R. 2006. Results of microbiological research in 2003 and 2004. EVI-EELA publications. Helsinki: Edita Prima Ltd. Pp. 29–36. (in Finnish)

Aarnisalo, K., Salo, S., Miettinen, H., Suihko, M.-L., Wirtanen, G., Autio, T., Lunden, J., Korkeala, H. and Sjöberg, A.-M. 2000. Bactericidal efficiencies of commercial disinfectants against *Listeria monocytogenes* on surfaces. J. Food Safety, Vol. 20, pp. 237–250.

Aase, B., Sundheim, G., Langsrud, S. and Rørvik, L. 2000. Occurrence of and possible mechanism for resistance to a quaternary ammonium compound in *Listeria monocytogenes*. Int. J. Food Microbiol., Vol. 62, pp. 57–63.

Aebi, R., Mühlemann, M., Bühlmann, G. and Schällibaum, M. 2003. Risk Assessment of *L. monocytogenes* in Swiss Emmental cheese. AgrarForschung, Vol. 10, pp. 306–311. (in German)

Allerberger, F. and Frischel, S. J. 1999. Use of automated ribotyping of Austrian *Listeria monocytogenes* isolates to support epidemiological typing. J. Microbiol. Meth., Vol. 35, p. 237–244.

Angelillo, I. F., Viggiani, N. M. A., Rizzo, L. and Bianco, A. 2000. Food handlers and foodborne diseases: Knowledge, attitudes and reported behaviour in Italy. J. Food Prot., Vol. 63, pp. 381–385.

Anonymous. 1989. Listeriosis from pre-cooked chicken reported in Britain. Food Chem. News, Vol. 30, pp. 26–27.

Anonymous. 1993. Hygienic equipment design criteria. Trends Food Sci. Tech., Vol. 4, pp. 225–229.

Anonymous. 1995. Hygienic design of equipment for open processing. Trends Food Sci. Tech., Vol. 6, pp. 305–310.

Anonymous. 1998. Directive 98/37/EC of the European Parliament and of the Council of 22 June 1998 on the approximation of the laws of the Member States relating to machinery as amended by Directive 98/79/EC. 23.7.98. OJ, L207, pp. 1–46.

Anonymous. 2000a. Outbreak of listeriosis linked to the consumption of rillettes in France. Eurosurveillance weekly 3, 19 Jan. 2000. Available at: http://www.eurosurv.org/update/news.htm.

Anonymous. 2000b. Zoonoses in Finland in 1995–1999. A report of the Ministry of Agriculture and Forestry 8/2000. Helsinki. 59 p.

Anonymous. 2003. Production and use of food-grade lubricants. EHEDG Update. Trends Food Sci. Tech., Vol. 14, pp. 157–162.

Anonymous. 2004a. Regulation (EC) No. 852/2004 of the European Parliament and of the Council of 29 April 2004 on the hygiene of foodstuffs. 25.6.2004. OJ, L226, pp. 3–21.

Anonymous. 2004b. Regulation (EC) No 1935/2004 of the European Parliament and of the Council of 27 October 2004 on materials and articles intended to come into contact with food and repealing Directives 80/590/EEC and 89/109/EEC. 13.11.2004. OJ, L338, pp. 4–17.

Anonymous. 2005. Regulation (EC) No. 2073/2005 on Microbiological Criteria for Foodstuffs. 22.12.2005. OJ, L338, pp. 1–26.

Anonymous. 2006. Regulation (EC) No. 2023/2006 of 22 December 2006 on good manufacturing practice for materials and articles intended to come into contact with food. 29.12.2006. OJ, L384, pp. 75–78.

Arimi, S. M., Ryser, E. T., Pritchard, T. J. and Donnelly, C. W. 1997. Diversity of *Listeria* ribotypes recovered from dairy cattle, silage, and dairy processing environments. J. Food Prot., Vol. 60, pp. 811–816.

Askarian, M., Kabir, G., Aminbaig, M., Memish, Z. A. and Jafari, P. 2004. Knowledge, attitudes, and practices of food service staff regarding food hygiene in Shiraz, Iran. Infect. Control Hosp. Epidemiol., Vol. 25, pp. 16–20.

Aureli, P., Fiorucci, G. C., Caroli, D., Marchiaro, G., Novara, O., Leone, L. and Salmaso, S. 2000. An outbreak of febrile gastroenteritis associated with corn contaminated by *Listeria monocytogenes*. New Engl. J. Med., Vol. 342, pp. 1236–1241.

Autio, T. 2003. Tracing the sources of *Listeria monocytogenes* contamination and listeriosis using molecular tools. Acad. Diss. Faculty of Vet. Med. Kuopio, Finland: Kopijyvä. 67 p. ISBN 952-91-6362-2.

Autio, T., Hielm, S., Miettinen, M., Sjöberg, A.-M., Aarnisalo, K., Björkroth, J., Mattila-Sandholm, T. and Korkeala, H. 1999. Sources of *Listeria monocytogenes* contamination in a cold-smoked rainbow trout processing plant detected by pulsed-field gel electrophoresis typing. Appl. Environ. Microbiol., Vol. 65, pp. 150–155.

Baloga, A. O. and Harlander, S. K. 1991. Comparison of methods for discrimination between strains of *Listeria monocytogenes* from epidemiological surveys. Appl. Environ. Microbiol., Vol. 57, pp. 2324–2331.

Bell, C. and Kyriakides, A. 2004. *Listeria monocytogenes*. In: Blackburn, C. de W. and McClure, P. J. (eds.). Foodborne pathogens: Hazards, risk analysis and control. Abington Hall, Abington, Cambridge: Woodhead Publishing Ltd. Pp. 337–361. ISBN 1-85573-454-0.

Bemrah, N., Cassin, M. H., Griffiths, M. W. and Cerf, O. 1998. Quantitative risk assessment of human listeriosis from consumption of soft cheese made from raw milk. Prev. Vet. Med., Vol. 37, pp. 129–145.

Bennett, R. W. 1986. Detection and quantitation of Gram-positive nonsporeforming pathogens and their toxins. In: Pierson, M. D. and Stern, N. J. (eds.). Foodborne microorganisms and their toxins: Developing methodology. USA: Marcel Dekker, Inc. Pp. 345–353. ISBN 0-8247-7607-0.

Berrang, M. E., Frank, J. F. and Brackett, R. E. 1988. Behaviour of *Listeria monocytogenes* in chocolate milk and ice cream mix made from post-expiration date skim milk. J. Food Prot., Vol. 51, p. 823.

Berrang, M. E., Meinersmann, R. J., Northcutt, J. K. and Smith, D. P. 2002. Molecular characterization of *Listeria monocytogenes* isolated from a poultry further processing facility and from fully cooked product. J. Food Prot., Vol. 65, pp. 1574–1579.

Best, M., Kennedy, M. E. and Coates, F. 1990. Efficacy of a variety of disinfectants against *Listeria* spp. Appl. Environ. Microbiol., Vol. 56, pp. 377–380.

Bille, J. 1990. Epidemiology of human listeriosis in Europe, with special reference to the Swiss outbreak. In: Miller, A. J., Smith, J. L. and Somkuti, G. A. (eds.). Foodborne Listeriosis. Amsterdam: Elsevier. Pp. 71–74.

Blackman, I. C. and Frank, J. F. 1996. Growth of *Listeria monocytogenes* as a biofilm on various food-processing surfaces. J. Food Prot., Vol. 59, pp. 827–831.

Brosch, R., Brett, M., Catimel, B., Luchansky, J. B., Ojeniyi, B. and Rocourt, J. 1996. Genomic fingerprinting of 80 strains from the WHO multicenter international typing study of *Listeria monocytogenes* via pulsed-field gel electrophoresis (PFGE). Int. J. Food Microbiol., Vol. 32, pp. 343–355.

Bruce, J. 1996. Automated system rapidly identifies and characterizes micro-organisms in food. Food Technol., Vol. 50, pp. 77–81.

Buchanan, R. L., Damert, W. G., Whiting, R. C. and van Schothorst, M. 1997. Use of epidemiological and food survey data to estimate a purposefully conservative dose-response relationship for *Listeria monocytogenes* levels and incidence of listeriosis. J. Food Prot., Vol 60, pp. 918–922.

Buchanan, R. L., Stahl, H. G. and Whiring, R. C. 1989. Effects and interactions of temperature, pH, atmosphere, sodium chloride and sodium nitrite on growth of *Listeria monocytogenes*. J. Food Prot., Vol. 52, pp. 844–851.

Carpenter, S. L. and Harrison, M. A. 1989. Survival of *Listeria monocytogenes* on processed poultry. J. Food Sci., Vol. 54, pp. 556–557.

CEN. 1997. EN 1672-2, 'Food processing machinery – Basic concepts – Part 2: Hygiene requirements'. Brussels, Belgium: European Committee for Standardization. 59 p.

Charaf, U. K., Bakich, S. L. and Falbo, D. M. 1999. A model biofilm for efficacy assessment of antimicrobials versus biofilm bacteria. In: Wimpenny, J., Gilbert, P., Walker, J., Brading, M. and Bayston, R. (eds.). Biofilms: The good, the bad and the ugly. Cardiff: BioLine. Pp. 171–177. ISBN 0-9520432-6-2.

Chen, Y., Ross, W. H., Scott, V. N. and Gombas, D. E. 2003. *Listeria monocytogenes*: Low levels equal low risk. J. Food Prot., Vol. 66, pp. 570–577.

Clayton, D. A., Griffith, C. J., Price, P. and Peters, A. C. 2002. Food handlers' beliefs and self-reported practices. Int. J. Environ. Health Res., Vol. 12, pp. 25–39.

Codex Alimentarius. 1999. Codex Alimentarius Comission. Principles and guidelines for the conduct of microbial risk assessment. CAC/GL-30. 6 p. [Accessed 2.7.2007.] Available at: http://www.codexalimentarius.net/download/standards/357/CXG 030e.pdf.

Cordoba, M. G., Cordoba, J. J. and Jordano, R. 1998. Evaluation of microbial hazards during processing of Spanish prepared flamenquin. J. Food Prot., Vol. 61, pp. 693–699.

Cox, N. A. and Bailey, J. S. 1999. Incidence and behaviour of *Listeria monocytogenes* in poultry and egg products In: Ryser, E. T. and Marth, E. H. (eds.). *Listeria*, Listeriosis, and Food Safety. 2nd ed. New York: Marcel Dekker. Pp. 565–600. ISBN 0-8247-0235-2.

Cummings, A. J., Fielding, A. K. and McLauchlin, J. 1994. Infection by *Listeria ivanovii* in a patient with AIDS. J. Infect., Vol. 28, pp. 89–91.

Curtis, G. D. W., Mitchell, R. G., King, A. F. and Griffin, E. J. 1989. Culture media and methods for the isolation of *Listeria monocytogenes*. Lett. Appl. Microbiol., Vol. 8, pp. 95-98.

Dalton, G. B., Austin, C. C., Sobel, J., Hayes, P. S., Bibb, W. F., Graves, L., Swaminathan, B., Proctor, M. E. and Griffin, P. M. 1997. An outbreak of gastroenteritis and fever due to *Listeria monocytogenes* in milk. New Engl. J. Med., Vol. 336, pp. 100–105.

Danchaivijitr, S., Rongrungruang, Y., Kachintorn, U., Techasathit, V., Pakaworavuthi, S. and Kachintorn, K. 2005. J. Med. Assoc. Thai., Vol. 88, Suppl. 10, pp. S31–S35.

Den Aantrekker, E. D., Boom, R. M., Zwietering, M. H. and van Schothorst, M. 2003. Quantifying recontamination through factory environments – review. Int. J. Food Microbiol., Vol. 80, pp. 117–130.

Destro, M. T., Leitao, M. F. and Farber, J. M. 1996. Use of molecular typing methods to trace the dissemination of *Listeria monocytogenes* in a shrimp processing plant. Appl. Environ. Microbiol., Vol. 62, pp. 705–711.

Dhaliwal, D. S., Cordier J. L. and Cox, L. J. 1992. Impedimetric evaluation of disinfectants against biofilms. Lett. Appl. Microbiol., Vol. 15, pp. 217–221.

Donnelly, C. W. and Baigent, G. J. 1986. Method for flow cytometric detection of *Listeria monocytogenes* in milk. Appl. Environ. Microbiol., Vol. 52, pp. 689–695.

Duncanson, P., Wareing, D. R. A. and Jones, O. 2003. Application of an automated immunomagnetic separation-enzyme immunoassay for the detection of *Salmonella* spp. during an outbreak associated with a retail premises. Lett. Appl. Microbiol., Vol. 37, pp. 144–148.

Earnshaw, A. M. and Lawrence, L. M. 1998. Sensitivity to commercial disinfectants, and the occurence of plasmids within various *Listeria monocytogenes* genotypes isolated from poultry products and the poultry processing environment. J. Appl. Microbiol., Vol. 84, pp. 642–648.

El-Kest, S. E. and Marth, E. H. 1988a. Inactivation of *Listeria monocytogenes* by chlorine. J. Food Prot. Vol. 51, pp. 520–524.

El-Kest, S. E. and Marth, E. H. 1988b. Temperature, pH and strain of pathogen as factors affecting inactivation of *Listeria monocytogenes* by chlorine. J. Food Prot., Vol. 51, pp. 622–625.

Embarek, P. K. B. and Huss, H. H. 1993. Heat resistance of *Listeria monocytogenes* in vacuum packaged pasteurized fish fillets. Int. J. Food Microbiol., Vol 20, pp. 85–95.

Ericsson, H., Eklöw, A., Danielsson-Tham, M.-L., Loncerevic, S., Mentzing, L.-O., Persson, I., Unnerstad, H. and Tham, W. 1997. An outbreak of listeriosis suspected to have been caused by rainbow trout. J. Clin. Microbiol., Vol. 35, pp. 2904–2907.

FAO/WHO. 2004. Risk assessment of *Listeria monocytogenes* in ready-to-eat foods. Microbiological risk assessment series; no. 5. [Accessed 8.8.2007.] Available at: http://www.who.int/foodsafety/publications/micro/en/mra5_contents.pdf.

Farber, J. M. (ed.). 1996. Special issue: Molecular typing of *Listeria*. Int. J. Food Microbiol., Vol. 32, pp. 251–366.

Farber, J. M. and Harwig, J. 1996. The Canadian position on *Listeria monocytogenes* in read-to-eat foods. Food Control, Vol. 7, pp. 253–258.

Farber, J. M. and Peterkin, P. I. 1991. *Listeria monocytogenes*, a food-borne pathogen. Microbiol. Rev., Vol. 55, pp. 476–511.

Farber, J. M., Coates, F. and Daley, E. 1992. Minimum water activity requirements for the growth of *Listeria monocytogenes*. Lett. Appl. Microbiol., Vol. 15, pp. 103–105.

Farber, J. M., Ross, W. H. and Harwig, J. 1996. Health risk assessment of *Listeria monocytogenes* in Canada. Int. J. Food Microbiol., Vol. 30, pp. 145–156.

FDA. 2000. Report of the FDA Retail Food Program Database of Foodborne Illness Risk Factors. US Food and Drug Administration, Center for Food Safety and Applied Nutrition, Washington DC, August 10, 2000. 59 p. [Accessed 8.8.2007.] Available at: http://www.cfsan.fda.gov/~acrobat/retrsk.pdf.

FDA/USDA. 2003. Quantitative assessment of the relative risk to public health from foodborne *Listeria monocytogenes* among selected categories of ready-to-eat foods. 541 p. [Accessed 8.8.2007.] Available at: http://www.foodsafety.gov/~dms/lmr2-toc.html.

Fleming, D. W., Cochi, S. L., MacDonald, K. L., Brondum, J., Hayes, P. S., Plikaytis, B. D., Holmes, M. B., Audurier, A., Broome, C. V. and Reincold, A. L. 1985. Pasteurized milk as a vehicle of infection in an outbreak of listeriosis. N. Engl. J. Med., Vol. 312, pp. 404–407.

Flores, R. A., Tamplin, M. L., Marmer, B. S., Phillips, J. G. and Cooke, P. H. 2006. Transfer coefficient models for *Escherichia coli* O157:H7 on contacts between beef tissue and high density polyethylene surfaces. J. Food Prot., Vol. 69, pp. 1248–1255.

Franco, C. M., Quinto, E. J., Fente, C., Rodriguez-Otero, J. L., Dominguez, L. and Cepeda, A. 1995. Determination of the principal sources of *Listeria* spp. contamination in poultry meat and a poultry processing plant. J. Food Prot., Vol. 58, pp. 1320–1325.

Frank, J. F. and Koffi, R. A. 1990. Surface-adherent growth of *Listeria monocytogenes* is associated with increased resistance to surfactant sanitizers and heat. J. Food Prot., Vol. 53, pp. 550–554.

Fraser, J. A. and Sperber, W. H. 1988. Rapid detection of *Listeria* spp. in food and environmental samples by esculin hydrolysis. J. Food Prot., Vol. 51, pp. 762–765.

Frye, D. M., Zweig, R., Sturgeon, J., Tormey, M., LeCavalier, M., Lee, I., Lawani, L. and Mascola, L. 2002. An outbreak of febrile gastroenteritis associated with delicatessen meat contaminated with *Listeria monocytogenes*. CID, Vol. 35, pp. 943–949.

FSIS. 2003. Draft FSIS risk assessment for *Listeria monocytogenes* in deli meats. 96 p. [Accessed 8.8.2007.] Available at http://www.fsis.usda.gov/OPPDE/rdad/FRPubs/97-013F/ListeriaReport.pdf.

Gendel, S. M. and Ulaszek, J. 2000. Ribotype analysis of strain distribution in *Listeria monocytogenes*. J. Food Prot., Vol. 63, pp. 179–185.

Golden, D. A., Beuchnat, L. R. and Brackett, R. E. 1988. Inactivation and injury of *Listeria monocytogenes* as affected by heating and freezing. Food Microbiol., Vol. 5, pp. 17–23.

Gottlieb, S. L., Newbern, E. C., Griffin, P. M., Graves, L. M., Hoekstra, R. M., Baker, N. L., Hunter, S. B., Holt, K. G., Ramsey, F., Head, M., Levine, P., Johnson, G., Schoonmaker-Bopp, D., Reddy, V., Kornstein, L., Gerwel, M., Nsubuga, J., Edwards, L., Stonecipher, S., Hurd, S., Austin, D., Jefferson, M. A., Young, S. D., Hise, K., Chernak, E. D., Sobel, J.; Listeriosis Outbreak Working Group. 2006. Multistate outbreak of listeriosis linked to turkey deli meat and subsequent changes in US regulatory policy. Clin. Infect. Dis. Vol. 42, pp. 29–36.

Goulet, V., Lepoutre, A., Rocourt, J., Courtier, A. L., Dehaumont, P. and Veit, P. 1993. Epidémie de listériose en France – bilan final et resultants de l'enquênte épidémiogique. Bull. Epidémiol. Hebdom., Vol. 4, pp. 13–14.

Goulet, V., Rocourt, J., Rebiere, I., Jacquet, C., Moyse, C., Dehaumont, P., Salvat, G. and Veit, P. 1998. Listeriosis outbreak associated with the consumption of rillettes in France in 1993. J. Infect. Dis., Vol. 177, pp. 155–160.

Gravani, R. 1999. Incidence and control of *Listeria* in food-processing facilities. In: Ryser, E. T. and Marth, E. H. (eds.). *Listeria*, Listeriosis, and Food Safety. 2nd ed. New York: Marcel Dekker. Pp. 657–709. ISBN 0-8247-0235-2.

Graves, L. M. and Swaminathan, B. 2001. PulseNet standardized protocol for subtyping *Listeria monocytogenes* by macrorestriction and pulse-field gel electrophoresis. Int. J. Food Microbiol., Vol. 65, pp. 55–62.

Graves, L. M., Swaminathan, B., Reeves, M. W., Hunter, S. B., Weaver, R. E., Plikaytis, B. D. and Schuchat, A. 1994. Comparison of ribotyping and multilocus enzyme electrophoresis for subtyping of Listeria monocytogenes isolates. J. Clin. Microbiol., Vol., 32, pp. 2936–2943.

Graves, L. M., Swaminathan, B. and Hunter, S. 1999. In: Ryser, E. T. and Marth, E. H. (eds.). *Listeria*, Listeriosis, and Food Safety. 2nd ed. New York: Marcel Dekker. Pp. 279–297. ISBN 0-8247-0235-2.

Gray, M. L. and Killinger, A. H. 1966. *Listeria monocytogenes* and listeric infections. Bacteriol. Rev., Vol. 30, pp. 309–382.

Grif, K., Heller, I., Wagner, M., Dierich, M. and Würzner, R. 2006. A comparison of *Listeria monocytogenes* serovar 4b isolates of clinical and food origin in Austria by automated ribotyping and pulsed-field gel electrophoresis. National and International PulseNet Programs. Foodborne Pathog Dis., Vol. 3, pp. 138–141.

Guzewich, J. and Ross, M. 1999. Evaluation of risks related to microbiological contamination of ready-to-eat food by food preparation workers and the effectiveness of interventions to minimize those risks. 28 p. [Accessed 15.11.2006.] Available at: http://www.cfsan.fda.gov/~ear/rterisk.html.

Haas, C. N. 1983. Estimation of the risk due to low doses of microorganisms: A comparison of alternative methodologies. Am. J. Epidemiol., Vol. 118, pp. 573–582.

Haas, C. N., Madabusi, A. T., Rose, J. B. and Gerba, C. P. 1999. Development and validation of dose response relationship from *Listeria monocytogenes*. Quant. Microbiol., Vol. 1, pp. 89–102.

Hamilton, W. A. 1991. Sulphate-reducing bacteria and their role in biocorrosion. In: Flemming, H.-C. and Geesey, G. G. (eds.). Biofouling and biocorrosion in industrial water systems. Berlin, Germany: Springer-Verlag. Pp. 187–193.

Hatakka, M. and Johansson, T. 2002. *Listeria* monocytogenes in vacuum-packed coldsmoked fish. (*Listeria monocytogenes* tyhjiöpakatuissa kylmäsavukaloissa.) Kaari, Vol. 1, No. 6, pp. 18–19. (in Finnish)

Hatakka, M. and Maijala, R. 2003. Official investigations of pathogenic bacteria in 2001. (Patogeenisten bakteerien viranomaistutkimukset vuonna 2001.) Kaari, Vol. 2, No. 2, pp. 26–27. (in Finnish)

Haupt, H., Köfer, J. and Fuchs, K. 1999. Abweichungsrate von Hygienevorschriften, Risikobewertung und Verbesserungsansätze in Frischfleisch-Klein- und Mittelunternehmen. Wien. Tierärtzl. Mschr., Vol. 86, pp. 122–128. (in German)

Health Canada. 2004. Policy on *Listeria monocytogenes* in ready-to-eat foods. Food Directorate–Health Products and Food Branch–Health Canada. Pp. 1–19.

Heinzel, M. 1988. The phenomena of resistance to disinfectants and preservatives. In: Payne, K. R. (ed.). Industrial biocides. Chichester, UK: John Wiley and Sons. Pp. 52–67.

Henroid, D. and Sneed, J. 2004. Readiness to implement hazard analysis and critical control point (HACCP) systems in Iowa schools. J. Am. Diet. Ass., Vol. 104, pp. 180–185.

Heuvelink, A. E., Roessink, G. L., Bosboom, K. and de Boer, E. 2001. Zero-tolerance for faecal contamination of carcasses as a tool in the control of O157 VTEC infections. Int. J. Food Microbiol., Vol. 66, pp. 13–20.

Hielm, S., Tuominen, P., Aarnisalo, K., Raaska, L. and Maijala, R. 2006. Attitudes towards own-checking and HACCP plans among Finnish food industry employees. Food Control, Vol. 17, pp. 402–407.

Holah, J. T. and Taylor, J. 2003. Personal hygiene. In: Lelieveld, H. L. M., Mostert, M. A., Holah, J. and White, B. (eds.). Hygiene in food processing. 1st ed. Cambridge: Woodhead Publishing limited. Pp. 288–309. ISBN 1-85573-466-4.

Hoornstra, E. and Notermans, S. 2001. Quantitative microbiological risk assessment. Int. J. Food Microbiol., Vol. 66, pp. 21–29.

Hoornstra, E., Northolt, M. D., Notermans, S. and Barendsz, A. W. 2001. The use of quantitative risk assessment in HACCP. Food Control, Vol. 12, pp. 229–234.

Hsu, J. C. 1991. Effective control of Listeria monocytogenes in a dairy processing and packaging plant by isothiazoline microbicide. Dairy Food Environ. Sanit., Vol. 11, pp. 70–72.

Hudson, J. A., Mott, S. J. and Penney, N. 1994. Growth of *Listeria monocytogenes*, *Aeromonas hydrophila*, and *Yersinia enterocolitica* on vacuum and saturated carbon dioxide controlled atmosphere-packaged sliced roast beef. J. Food Prot., Vol. 57, pp. 204–208.

Humphrey, T. J. and Worthington, D. M. 1990. *Listeria* contamination of retail meat slicers. PHLS Microbiology Digest, Vol. 7, p. 57.

Hunter, P. R. and Gaston, M. A. 1988. Numerical index of the discriminatory ability of typing systems: An application of Simpson's index of diversity. J. Clin. Microbiol., Vol. 26, pp. 2465–2466.

Husu, J., Johansson, T., Oivanen, L. and Hirn, J. 1993. *Listeria monocytogenes* in poultry and poultry meat in Finland. In: Nurmi, E., Colin, P. and Mulder, R. W. A. W. (eds.). Other pathogens of concern (no Salmonella and Campylobacter), FLAIR No. 6/COST No. 106, Proceedings 8, Meeting held at Helsinki, Finland, June 11–14th, 1992. Pp. 85–91.

ICMSF. 1996. Roberts, T. A., Baird-Parker, A. C. and Tompkin, R. B. (eds.). Microorganisms in foods 5, Characteristics of microbial pathogens. 1st ed. Suffolk, Great Britain: St. Edmundsbury Press Ltd. Pp. 141–182. ISBN 0-412-47350-X.

ISO. 1996. Microbiology of food and animal feeding stuffs – Horizontal method for the detection and enumeration of *Listeria monocytogenes* – Part 1: Detection method, ISO 11290-1. Geneva, Switzerland: International Organisation for Standardization. 20 p.

ISO. 1998. Microbiology of food and animal feeding stuffs – Horizontal method for the detection and enumeration of *Listeria monocytogenes* – Part 2: Enumeration method. ISO 11290-2. Geneva, Switzerland: International Organisation for Standardization. 20 p.

ISO. 2002. Safety of machinery – Hygiene requirements for the design of machinery. ISO 14159:2002. Geneva, Switzerland: International Organisation for Standardization. 20 p.

ISO. 2006. Safety of machinery – Lubricants with incidental product contact – Hygiene requirements. ISO 21469:2006. Geneva, Switzerland: International Organisation for Standardization. 7 p.

Ivanek, R., Gröhn, Y. T., Wiedmann, M. and Wells, M. T. 2004. Mathematical model of *Listeria monocytogenes* cross-contamination in a fish processing plant. J. Food Prot., Vol. 67, pp. 2688–2697.

Jay, J. M. 1996. Modern food microbiology. 5th ed. New York: Chapman & Hall. Pp. 478–506. 661 p. ISBN 0-412-07691-8.

Jensen, N. E., Aarestrup, F. M., Jensen, J. and Wegener, H. C. 1996. *Listeria monocytogenes* in bovine mastitis. Possible implication for human health. Int. J. Food Microbiol., Vol. 32, pp. 209–216.

Jeong, D. K. and Frank, J. F. 1994. Growth of *Listeria monocytogenes* at 10 °C in biofilms with microorganisms isolated from meat and dairy processing environments. J. Food Prot., Vol. 57, pp. 576–586.

Johansson, T. 1998. Enhanced detection and enumeration of *Listeria monocytogenes* from foodstuffs and food-processing environments. Int. J. Food Microbiol., Vol. 40, pp. 77–85.

Johansson, T. and Hirn, J. 1992. Comparison of DNA-hybridization, ELISA and culturing methods for detection of *Salmonella* and *Listeria* in deep-frozen broiler carcasses. In: Schwarzwer, Chr. and Mulder, R. W. A. W. (eds.). Detection methods and sampling plans for pathogens in poultry, FLAIR No. 6/COST No. 106, Proceedings 6. Meeting held at Grub, Germany, September 22–24th, 1991. Pp. 71–78.

Johansson, T., Rantala, L., Palmu, L. and Honkanen-Buzalski, T. 1999. Occurrence and typing of *Listeria monocytogenes* strains in retail vacuum-packed fish products and in a production plant. Int. J. Food Microbiol., Vol. 47, pp. 111–119.

Johansson, T., Hatakka, M. and Maijala, R. 2002. Summary from results of a research project on cold cuts (Yhteenveto tutkimusprojektin tuloksista lihaleikkeleet EELA-EVI 2002). National Food Agency (EVI) and National Veterinary and Food Research Institute (EELA). 5 p. [Accessed 8.8.2007.] Available at: http://www.evira.fi/attachments/ elintarvikkeet/valvonta_ja_yrittajat/tutkimukset_ja_projektit/arkisto/lihaleikkeleet-tutkimusprojektin_yhteenveto.pdf. (in Finnish)

Johansson, T., Markkula, A., Hatakka, M., Oivanen, L. and Maijala, R. 2003. Manual of microbiological hazards connected with food and drinking water. Publication 1/2003. Helsinki: National Food Agency (EVI) and National Veterinary and Food Research Institute (EELA). 211 p. ISBN 951-732-185-6. (in Finnish)

Junttila, J. and Brander, M. 1989. *Listeria monocytogenes* septicemia associated with consumption of salted mushrooms. Scand. J. Inf. Dis., Vol. 21, pp. 339–342.

Kaczmarski, E. B. and Jones, D. M. 1989. Listeriosis and ready-cooked chicken. Lancet, Vol. i, No. 8637, p. 549.

Kerouanton, A., Brisabois, A., Denoyer, E., Dilasser, F., Grout, J., Salvat, G. and Picard, B. 1998. Comparison of five typing methods for the epidemiological study of *Listeria monocytogenes*. Int. J. Food Microbiol., Vol. 43, pp. 61–71.

Khelef, N., Lecuit, M., Buchrieser, C., Cabanes, D., Dussurget, O. and Cossart, P. 2006. *Listeria monocytogenes* and the genus *Listeria*. In: Dworkin, M. (ed.). The prokaryotes: An evolving electronic resource for the microbiological community. [Accessed 22.9.2006.] Available at: http://141.150.157.117:8080/prokPUB/.

Kirby, M. P. and Gardiner, K. 1997. The effectiveness of hygiene training for food handlers. Int. J. Environ. Health. Res., Vol. 7, pp. 251–258.

Kleer, J. and Hildebrandt, G. 2001. Bedeutung der Predictive Microbiology zur Risiko-Minimierung im Herstellungsprozess. Fleischwirtsch., Vol. 81, No. 6, pp. 99–2001. Krysinski, E. P., Brown, L. J. and Marchisello T. J. 1992. Effect of cleaners and sanitizers on *Listeria monocytogenes* attached to product contact surfaces. J. Food Prot., Vol. 55, pp. 246–251.

Köhler, C. R. 2001. Food-grade lubricants for the food processing industry. New Food, Vol. 4, pp. 49–54.

LaPorte, T., Heisey-Grove, D., Kludt, P., Matyas, B. T., De Maria Jr., A., Dicker, R., De, A., Fiore, A., Nainan, O. and Friedman, D. S. 2003. Foodborne transmission of hepatitis A – Massachusetts. MMWR, Vol. 52, pp. 565–567.

Lee, S.-H. and Frank, J. F. 1991a. Effect of growth temperature and media on inactivation of *Listeria monocytogenes* by chlorine. J. Food Safety, Vol. 11, pp. 65–71.

Lee, S.-H. and Frank, J. F. 1991b. Inactivation of surface-adherent Listeria monocytogenes. Hypochlorite and heat. J. Food Prot., Vol. 54, pp. 4–6.

Lee, W. H. and McClain, D. 1986. Improved *Listeria monocytogenes* selective agar. Appl. Environ. Microbiol., Vol. 52, pp. 1215–1217.

Lelieveld, H. L. M., Mostert, M. A. and Curiel, G. J. 2003. Hygienic equipment design. In: Lelieveld, H. L. M., Mostert, M. A., Holah, J. and White, B. (eds.). Hygiene in food processing. 1st ed. Cambridge: Woodhead Publishing limited. Pp. 122–166. ISBN 1-85573-466-4.

Lessing, M. P. A., Curtis, G. D. W. and Bowler, I. C. J. 1994. *Listeria ivanovii* infection. J. Infect., Vol. 29, pp. 230–231.

Lewan, M. 2003. Equipment construction materials and lubricants In: Lelieveld, H. L. M., Mostert, M. A., Holah, J. and White, B. (eds.). Hygiene in food processing. 1st ed. Cambridge: Woodhead Publishing limited. Pp. 167–178. ISBN 1-85573-466-4.

Lin, C. M., Takeuchi, K., Zhang, L., Dohm, C. B., Meyer, J. D., Hall, P. A. and Doyle, M. P. 2006. Cross-contamination between processing equipment and deli meats by *Listeria monocytogenes*. J. Food Prot., Vol. 69, pp. 71–79.

Lindqvist, R. and Westöö, A. 2000. Quantitative risk assessment for Listeria monocytogenes in smoked or gravad salmon and rainbow trout in Sweden. Int. J. Food Microbiol., Vol. 58, pp. 181–196.

Linnan, M. J., Mascola, L., Lou, X. D., Goulet, V., May, S., Salminen, C., Hird, D. W., Yonekura, M. L., Hayes, P., Weaver, R., Audurier, A., Plikaytis, B. D., Fannin, S. L., Kleks, A. and Broome, C. V. 1988. Epidemic listeriosis associated with Mexican-style cheese. New Engl. J. Med., Vol. 319, pp. 823–828.

Lou, Y. and Yousef, A. E. 1999. Characteristics of *Listeria monocytogenes* important to food processors. In: Ryser, T. Y. and Marth, E. H. (eds.). *Listeria*, Listeriosis and Food Safety. 2nd ed. New York: Marcel Dekker, Inc. Pp. 131–224.

Louie, M., Jayaratne, P., Luchsinger, I., Devenish, J., Yao, J., Schlech, W. and Simor, A. 1996. Comparison of ribotyping, arbitrarily primed PCR, and pulsed-field gel electrophoresis for molecular typing of *Listeria monocytogenes*. J. Clin. Microbiol., Vol. 34, pp. 15–19.

Lukinmaa, S., Aarnisalo, K., Suihko, M. L. and Siitonen, A. 2004. Diversity of *Listeria monocytogenes* isolates of human and food origin studied by serotyping, automated ribotyping and pulsed-field gel electrophoresis. Clin. Microbiol. and Infect., Vol. 10, pp. 562–568.

Lundén, J. 2004. Persistent *Listeria monocytogenes* contamination in food processing plants. Acad. Diss. Faculty of Vet. Helsinki, Finland: Yliopistopaino. 60 p. ISBN 952-91-6697-4.

Lundén, J. M., Miettinen, M. K., Autio, T. J. and Korkeala, H. J. 2000. Persistent *Listeria monocytogenes* strains show enhanced adherence to food contact surface after short contact times. J. Food Prot., Vol. 63, pp. 1204–1207.

Lundén, J. M., Autio, T. J. and Korkeala, H. J. 2002. Transfer of persistent *Listeria monocytogenes* contamination between food-processing plants associated with a dicing machine. J. Food Prot., Vol. 65, pp. 1129–1133.

Lundén, J., Autio, T., Markkula, A., Hellström, S. and Korkeala, H. 2003. Adaptive and cross-adaptive responses of persistent and non-persistent *Listeria monocytogenes* strains to disinfectants. Int. J. Food Microbiol., Vol. 82, pp. 265–272.

Lyhs, U., Hatakka, M., Mäki-Petäys, N., Hyytiä, E. and Korkeala, H. 1998. Microbiological quality of Finnish vacuum-packaged fishery products at retail level. Arch. Lebensmittelhyg., Vol. 49, pp. 146–150.

Lyytikäinen, O., Autio, T., Maijala, R., Ruutu, P., Honkanen-Buzalski, T., Miettinen, M., Hatakka, M., Mikkola, J., Anttila, V.-J., Johansson, T., Rantala, L., Aalto, T., Korkeala, H. and Siitonen, A. 2000. An outbreak of *Listeria monocytogenes* serotype 3a infections from butter in Finland. J. Infect. Dis., Vol. 181, pp. 1835–1841.

Lyytikäinen, O., Nakari, U. M., Lukinmaa, S., Kela, E., Minh, N. N. T. and Siitonen, A. 2006. Surveillance of listeriosis in Finland during 1995–2004. Euro Surveill., Vol. 11, pp. 82–85.

Mafu, A. A., Roy, D., Goulet, J. and Magny, P. 1990. Attachment of *Listeria monocytogenes* to stainless steel, glass, polyprophylene, and rubber surfaces after short contact times. J. Food Prot., Vol. 53, pp. 742–746.

Mai, T. L., Sofyan, N. I., Fergus, J. W., Gale, W. F. and Conner, D. E. 2006. Attachment of *Listeria monocytogenes* to an austenitic stainless steel after welding and accelerated corrosion treatments. J. Food Prot., Vol. 69, pp. 1527–1532.

Maijala, R., Lyytikäinen, O., Johansson, T., Autio, T., Haavisto, L. and Honkanen-Buzalski, T. 2001. Exposure of *Listeria monocytogenes* within an epidemic caused by butter in Finland. Int. J. Food Microbiol., Vol. 70, pp. 97–109.

Marriott, N. G. 1999. Principles of food sanitation. 4th ed. Gaithersburg, MD: Aspen Publishers, Inc. Pp. 60–74, 139–157. ISBN 0-8342-1232-3.

Martinez-Gonzales, N. E., Hernandez-Herrera, A., Martinez-Chavez, L., Rodriguez-Garcia, M. O., Torres-Vitela, M. R., Mota-de la Garza, L. and Castillo, A. 2003. Spread of bacterial pathogens during preparation of freshly squeezed orange juice. J. Food Prot., Vol. 66, pp. 1490–1494.

McDonnell, G. and Russell, D. 1999. Antiseptics and disinfectants: Activity, action, and resistance. Clin. Microbiol. Rev., Vol. 12, pp. 147–179.

McLauchlin, J., Hall, S. M., Velani, S. K., and Gilbert, R. J. 1991. Human listeriosis and pate: a possible association. Br. Med. J., Vol. 303, pp. 773–775.

McLauchlin, J., Mitchell, R. T., Smerdon, W. J. and Jewell, K. 2004. *Listeria monocytogenes* and listeriosis: a review of hazard characterization for use in microbiological assessment of foods. Int. J. Food Microbiol., Vol. 92, pp. 15–33.

McMeekin, T. A., Baranyi, J., Bowman, J., Dalgaard, P., Kirk, M., Ross, T., Schmid, S. and Zwietering, M. H. 2006. Information systems in food safety management. Int. J. Food Microbiol., Vol. 112, pp. 181–194.

Mead, G. C., Hudson, W. R. and Ariffin, R. 1990. Survival and growth of *Listeria* monocytogenes on irradiated poultry carcasses. Lancet, Vol. 335, No. 8696, p. 1036.

Michaels, B., Keller, C., Blevins, M., Paoli, G., Ruthman, T., Todd, E. and Griffith, C. J. 2004. Prevention of food worker transmission of foodborne pathogens: risk assessment and evaluation of effective hygiene intervention strategies. Food Service Tech., Vol. 4, pp. 31–49.

Midelet, G. and Carpentier, B. 2002. Transfer of micro-organisms, including *Listeria monocytogenes*, from various materials to beef. Appl. Environ. Microbiol., Vol. 68, pp. 4015–4024.

Midelet, G., Kobilinsky, A. and Carpentier, B. 2006. Construction and analysis of fractional multifactorial designs to study attachment strength and transfer of *Listeria monocytogenes* from pure or mixed biofilms after contact with a solid model food. Appl. Environ. Microbiol., Vol. 72, pp. 2313–2321.

Miettinen, H. and Wirtanen, G. 2006. Ecology of *Listeria* spp. in a fish farm and molecular typing of *L. monocytogenes* from fish farming and processing companies. Int. J. Food Microbiol., Vol. 112, pp. 138–146.

Miettinen, M. K., Björkroth, K. J. and Korkeala, H. J. 1999a. Characterization of *Listeria monocytogenes* from an ice cream plant by serotyping and pulsed-field gel electrophoresis. Int. J. Food Microbiol., Vol. 46, pp. 187–192.

Miettinen, M. K., Siitonen, A., Heiskanen, P., Haajanen, H., Björkroth, K. J. and Korkeala, H. J. 1999b. Molecular epidemiology of an outbreak of febrile gastroenteritis caused by *Listeria monocytogenes* in cold-smoked rainbow trout. J. Clin. Microbiol., Vol. 37, pp. 2358–2360.

Miettinen, M. K., Palmu, L., Björkroth, K. J. and Korkeala, H. 2001. Prevalence of *Listeria monocytogenes* in broilers at the abattoir, processing plant, and retail level. J. Food Prot., Vol. 64, pp. 994–999.

Miller, A. J., Whiting, R. C. and Smith, J. L. 1997. Use of risk assessment to reduce listeriosis incidence. Food Technol., Vol. 51, pp. 100–103.

Misrachi, A., Watson, A. J. and Coleman, D. 1991. *Listeria* in smoked mussels in Tasmania. Comm. Dis. Intell., Vol. 15, p. 427.

Mosteller, T. M. and Bishop, J. R. 1993. Sanitizer efficacy against attached bacteria in a milk biofilm. J. Food Prot., Vol. 56, pp. 34–41.

Murphy, R. Y., Osaili, T., Duncan, L. K. and Marcy, J. A. 2004. Thermal inactivation of Salmonella and Listeria monocytogenes in ground chicken thigh/leg meat and skin. Poultry Sci., Vol. 83, pp. 1218–1225.

Murray, E. G. D., Webb, R. A. and Swann, M. B. R. 1926. A disease of rabbits characterized by large mononuclear leucocytosis, caused by a hitherto undescribed bacillus, *Bacterium monocytogenes* (n. sp.). J. Pathol. Bacteriol., Vol. 29, pp. 407–439.

Mustapha, A. and Liewen, M. B. 1989. Destruction of *Listeria monocytogenes* by sodium hypochlorite and quaternary ammonium sanitizers. J. Food Prot., Vol. 52, pp. 306–311.

Møretrø, T. and Langsrud, S. 2004. Review article. *Listeria monocytogenes*: biofilm formation and persistence in food-processing environments. Biofilms, Vol. 1, pp. 107–121.

Nel, S., Lues, J. F. R., Buys, E. M. and Venter, P. 2004. The personal and general hygiene practices in the deboning room of a high throughput red meat abattoir. Food Control, Vol. 15, pp. 571–578.

Netuschil, K. 1995. Hygienic lubrication in the food processing and pharmaceutical industry. Food Tech Europe. Vol. 2, pp. 82, 84–86, 88–90.

NFA. 1998, 2000–2005. Official communication. Dairy plants own checks. Number of annual tests and positive results in years 1998, 2000–2005. Milk and Egg hygiene Unit. Helsinki: National Food Agency.

Norton, D. M., McCamey, M. A., Gall, K., Scarlett, J. M., Boor K. J. and Wiedmann, M. 2001. Molecular studies on the ecology of *Listeria monocytogenes* in the smoked fish processing industry. Appl. Environ. Microbiol., Vol. 67, pp. 198–205.

Norwood, D. E. and Gilmour, A. 1999. Adherence of *Listeria monocytogenes* strains to stainless steel coupons. J. Appl. Microbiol., Vol. 86, pp. 576–582.

Notermans, S. and Mead, G. C. 1996. Incorporation of elements of quantitative risk analysis in the HACCP system. Int. J. Food Microbiol., Vol. 30, pp. 157–173.

Ojeniyi, B., Wegener, H. C., Jensen, N. E. and Bisgaard, M. 1996. *Listeria monocytogenes* in poultry and poultry products: epidemiological investigations in seven Danish abattoirs. J. Appl. Bacteriol., Vol. 80, pp. 395–401.

Olsen, J. A., Yousef, A. E. and Marth, E. H. 1988. Growth and survival of *Listeria monocytogenes* during making and storage of butter. Milchwissenschaft, Vol. 43, pp. 487–489.

Olsen, S. J., Patrick, M., Hunter, S. B., Reddy, V., Kornstein, L., MacKenzie, W. R., Lane, K., Bidol, S., Stoltman, G. A., Frye, D. M., Lee, I., Hurd, S., Jones, T. F., LaPorte, T. N., Dewitt, W., Graves, L., Wiedmann, M., Schoonmaker-Bopp, D. J., Huang, A. J., Vincent, C., Bugenhagen, A., Corby, J., Carloni, E. R., Holcomb, M. E., Woron, R. F., Zansky, S. M., Dowdle, G., Smith, F., Ahrabi-Fard, S., Ong, A. R., Tucker, N., Hynes, N. A. and Mead, P. 2005. Multistate outbreak of *Listeria monocytogenes* infection linked to delicatessen turkey meat. Clin. Infect. Dis., Vol. 40, pp. 962–967.

Orth, R. and Mrozek, H. 1990. Is the control of *Listeria*, *Campylobacter* and *Yersinia* a disinfection problem? Fleischwirtsch., Vol. 69, pp. 1575–1576.

Ortiz, C., Guiamet, P. S. and Videla, H. A. 1990. Relationship between biofilms and corrosion of steel by microbial contaminants of cutting-oil emulsions. International Biodeterioration, Vol. 26, pp. 315–326.

Paulson, D. S. 2000. Handwashing, gloving and disease transmission by the food preparer. Dairy, Food and Environmental Sanitation, Vol. 20, pp. 838–845.

Pérez-Rodríguez, F., Todd, E. C. D., Valero, A., Carrasco, E., García, R. M. and Zurera, G. 2006. Linking quantitative exposure assessment and risk management using the food safety objective concept: An example with *Listeria monocytogenes* in different cross-contamination scenarios. J. Food Prot., Vol. 69, pp. 2384–2394.

Petitdemange, E., Durr, C., Andaloussi, S. A. and Raval, G. 1995. Fermentation of raw glycerol to 1,3-propanediol by new strains of *Clostridium butyricum*. J. Indust. Microbiol., Vol. 15, pp. 498–502.

Rahkio, M. and Korkeala, H. 1996. Microbiological contamination of carcasses related to hygiene practice and facilities on slaughtering lines. Acta Vet. Scand., Vol. 37, pp. 219–228.

Rasmussen, S. K. J., Ross, T., Olley, J. and McMeekin, T. 2002. A process risk model for the shelf life of Atlantic salmon fillets. Int. J. Food Microbiol., Vol. 73, pp. 47–60.

Reij, M. W. and den Aantrekker, E. D. 2004. Recontamination as a source of pathogens in processed foods. ILSI Europe Risk Analysis in Microbiology Task Force. Int. J. Food Microbiol., Vol. 91, pp. 1–11.

Reimer, L., Mottice, S. and Andrews, D. 1988. The effect of pH on survival of *Listeria monocytogenes*. Proc. Annual Meeting of the American Society for Microbiology, Miami Beach, May 8–13, Abstr. C-175.

Rocourt, J. 1996. Risk factors for listeriosis. Food Control, Vol. 7, pp. 195-202.

Rocourt, J., BenEmbarek, P., Toyofuku, H. and Schlundt, J. 2003. Quantitative risk assessment of *Listeria monocytogenes* in ready-to-eat foods: the FAO/WHO approach. FEMS Immunology & Medical Microbiology, Vol. 35, pp. 263–267.

Rodríguez, A. and McLandsborough, L. A. 2007. Evaluation of the transfer of *Listeria monocytogenes* from stainless steel and high-density polyethylene to bologna and American cheese. J. Food Prot., Vol. 70, pp. 600–606.

Ronner, A. B. and Wong, A. C. L. 1993. Biofilm development and sanitizer inactivation of *Listeria monocytogenes* and *Salmonella typhimurium* on stainless steel and Buna-n rubber. J. Food Prot., Vol. 56, pp. 750–758.

Rose, J. B., Haas, C. N. and Regli, S. 1991. Risk assessment and control of waterborne giardiasis. Am. J. Publ. Health, Vol. 88, pp. 709–713.

Ross, T. and McMeekin, T. A. 2003. Modeling microbial growth within food safety risk assessments. Risk Anal., Vol. 23, pp. 179–197.

Ross, T. and Sumner, J. 2002. A simple, spreadsheet-based, food safety risk assessment tool. Int. J. Food Microbiol., Vol. 77, pp. 39–53.

Rossmoore, K. 1988. The microbial activity of glutaraldehyde in chain conveyer lubricant formulations. In: Houghton, D. R., Smith, R. N. and Eggins, H. O. W. (eds.). Biodeterioration, Vol. 7. London: Elsevier Publishers Ltd. Pp. 242–247.

Ryser, E. T., Arimi, S. M., Bunduki, M. M.-C. and Donnelly, C. W. 1996. Recovery of different *Listeria* ribotypes from naturally contaminated, raw refrigerated meat and poultry products with two primary enrichment media. Appl. Environ. Microbiol., Vol. 62, pp. 1781–1787.

Rørvik, L. M. and Yndestad, M. 1991. *Listeria monocytogenes* in foods in Norway. Int. J. Food Microbiol., Vol. 13, pp. 97–104.

Sallam, S. S. and Donnelly, C. W. 1992. Destruction, injury, and repair of *Listeria* species exposed to sanitizing compounds. J. Food Prot., Vol. 55, pp. 771–776.

Samelis, J. and Metaxopoulos, J. 1999. Incidence and principal sources of *Listeria* spp. and *Listeria monocytogenes* contamination in processed meats and a meat processing plant. Food Microbiol., Vol. 16, pp. 465–477.

Samelis, J., Kakouri, A., Georgiadou, K. G. and Metaxopoulos, J. 1998. Evaluation of the extent and type of bacterial contamination at different stages of processing of cooked ham. J. Appl. Microbiol., Vol. 84, pp. 649–660.

Sanaa, M., Coroller, L. and Cerf, O. 2004. Risk assessment of listeriosis linked to the consumption of two soft cheeses made from raw milk: Camembert of Normandy and Brie of Meaux. Risk Analysis, Vol. 24, pp. 389–399.

Sasahara, K. C. and Zottola, E. A. 1993. Biofilm formation by *Listeria monocytogenes* utilizes a primary colonizing microorganism in flowing systems. J. Food Prot., Vol. 56, pp. 1022–1028.

Saunders, B. D., Schukken, Y., Kornstein, L., Reddy, V., Bannerman, T., Salehi, E., Dumas, N., Anderson, B. J., Massey, J. P. and Wiedmann, M. 2006. Molecular epidemiology and cluster analysis of human listeriosis cases in three U.S. States. J. Food Prot., Vol. 69, pp. 1680–1689.

Schaffner, D. W. 2004. Mathematical frameworks for modeling *Listeria* cross-contamination in food-processing plants. JFS, Vol. 69, pp. 155–159.

Schlech, W. F., Lavigne, P. M., Bortolussi, R. A., Allen, A. C., Haldane, E. V., Wort, A. J., Hightower, A. W., Johnson, S. E., King, S. H., Nicholls, E. S. and Broome, C. V. 1983. Epidemic listeriosis – evidence for transmission by food. N. Engl. J. Med., Vol. 308, pp. 203–206.

Schwartz, B., Ciesilski, C. A., Brome, C. V., Gaventa, S., Brown, G. R., Gellin, B. G., Hightower, A. W. and Mascola, L., the Listeriosis Study Group. 1988. Association of

sporadic listeriosis with consumption of uncooked hot dogs and undercooked chicken. Lancet, Vol. ii, No. 8614, pp. 779–782.

Seeliger, H. P. R. and Jones, D. 1986. Genus *Listeria* Pirie 1940, 383^{AL}. In: Sneath, P. H. A., Mair, N. S., Sharpe, M. E. and Holt, J. G. (eds.). Bergey's manual of Systematic Bacteriology. 2nd ed. Baltimore, MD: Williams and Wilkins. Pp. 1235–1245.

Serra, J. A., Domenach, E., Escriche, I. and Martorell, S. 1999. Risk assessment and critical control points from the production perspective. Int. J. Food Microbiol., Vol. 46, pp. 9–26.

Shin-Ho-Lee and Frank, J. F. 1991. Inactivation of surface-adherent *Listeria monocytogenes*, hypochlorite and heat. J. Food Prot., Vol. 54, pp. 4–6.

Skinner, R. 1996. *Listeria*: the state of the science, Rome 29–30 June 1995, Session IV: country and organizational postures on *Listeria monocytogenes* in food, *Listeria*: UK government's approach. Food Control, Vol. 7, pp. 245–247.

Slutsker, L. and Schuchat, A. 1999. Listeriosis in humans. In: Ryser, E. T. and Marth, E. H. (eds.). *Listeria*, Listeriosis, and Food Safety. 2nd ed. New York: Marcel Dekker. Pp. 74–95. ISBN 0-8247-0235-2.

Smoot, L. M. and Pierson, M. D. 1998. Effect of environmental stress on the ability of *Listeria monocytogenes* Scott A to attach to food contact surfaces. J. Food Prot., Vol. 61, pp. 1293–1298.

Sneed, J., Strohbehn, C. and Gilmore, S. A. 2004. Food safety practices and readiness to implement HACCP programs in assisted-living facilities in Iowa. J. Am. Diet Assoc., Vol. 104, pp. 1678–1683.

Suihko, M. L., Salo, S., Niclasen, O., Gudbjörnsdóttir, B., Torkelsson, G., Bredholt, S., Sjöberg, A. M. and Gustavsson, P. 2002. Characterization of *Listeria monocytogenes* isolates from the meat, poultry and seafood industries by automated ribotyping. Int. J. Food Microbiol., Vol. 72, pp. 137–146.

Sumner, J. and Ross, T. 2002. A semi-quantitative seafood safety risk assessment. Int. J. Food Microbiol., Vol. 77, pp. 55–59.

Swaminathan, B., Gerner-Smidt, P., Lai-King, Ng., Lukinmaa, S., Kai-Man, K., Rolando, S., Perez-Gutierrez, E. and Binsztein, N. 2006. Building PulseNet International: an interconnected system of laboratory networks to facilitate timely public health recognition and response to foodborne disease outbreaks and emerging foodborne diseases. Foodborne Pathog. Dis., Vol. 3, pp. 36–50.

Tang, C. H. and Fong, U. W. 2004. A survey of food hygiene knowledge and attitudes among Chinese food handlers in Fong Song Tong district. Asia Pac. J. Public Health, Vol. 16, pp. 121–125.

Taormina, P. J. and Beuchat, L. R. 2001. Survival and heat resistance of *Listeria monocytogenes* after exposure to alkali and chlorine. Appl. Environ. Microbiol., Vol. 67, pp. 2555–2563.

Tasara, T. and Stephan, R. 2006. Cold stress tolerance of *Listeria monocytogenes*: A review of molecular adaptive mechanisms and food safety implications. J. Food Prot., Vol. 69, pp. 1473–1484.

Teodorovic, R., Bugarski, M., Teodorovic, V., Radenkovic, B. and Mirilovic, M. 2000. Effect of disinfectants on different strains of *Listeria monocytogenes*. Acta Veterinaria (Beograd), Vol. 50, pp. 43–50.

Todd, E. C. D. and Harwig, J. 1996. Microbial risk analysis of food in Canada. J. Food Prot., 1996 Suppl., pp. 10–18.

Tompkin, R. B. 2002. Control of *Listeria monocytogenes* in the food-processing environment. J. Food Prot., Vol. 65, pp. 709–725.

Troller, J. A. 1993. Sanitation in food processing. 2nd ed. San Diego: Academic Press. ISBN 0-12-700655-9.

Tuncan, E. U. 1993. Effect of cold temperature on germicidal efficacy of quaternary ammonium compound, iodofor and chlorine on *Listeria*. J. Food Prot., Vol. 56, pp. 1029–1033.

Tuominen, P., Hielm, S., Aarnisalo, K., Raaska, R. and Maijala, R. 2003. Trapping the food safety performance of a small or medium-sized food company using a risk-based model. The HYGRAM[®] system. Food Control, Vol. 14, pp. 573–578.

Upmann, M. and Reuter, G. 1998. Oberflächenkeimgehalte und Betriebshygiene in einem Zerlegebetrieb für Schweinefleisch. Fleischwirtsch., Vol. 78, pp. 647–651.

Walker, S. J., Archer, P. and Banks, J. G. 1990. Growth of *Listeria monocytogenes* at refrigeration temperatures. J. Appl. Bacteriol., Vol. 68, pp. 157–162.

Walls, I. and Scott, V. N. 1997. Use of predictive microbiology in microbial food safety risk assessment. Int. J. Food Microbiol., Vol. 36, pp. 97–102.

van de Weyer, A., Devleeschouwer, M.-J. and Dony, J. 1993. Bactericidal activity of disinfectants on *Listeria*. J. Appl. Bacteriol., Vol. 74, pp. 480–483.

van der Waa, P. K. 1995. Lubrication of food processing machinery. In: Food Technology International Europe. London: Sterling Publications Ltd. Pp. 140–144.

van Gerwen, S. J. C. and Zwietering, M. H. 1998. Growth and inactivation models to be used in quantitative risk assessments. J. Food Prot., Vol. 61, pp. 1541–1549.

van Gerwen, S. J. C, te Giffel, M. C., van't Riet, K., Beumer, R. R. and Zwietering, M. H. 2000. Stepwise quantitative risk assessment as a tool for characterization of microbiological food safety. J. Appl. Microbiol., Vol. 88, pp. 938–951.

van Netten, P., Perales, I., van de Moosdijk, A., Curtis, G. D. W. and Mossel, D. A. A. 1989. Liquid and solid selective differential media for the detection and enumeration of *L. monocytogenes* and other *Listeria* spp. Int. J. Food Microbiol., Vol. 8, pp. 299–316.

van Schothorst, M. 1997. Practical approaches to risk assessment. J. Food Prot., Vol. 60, pp. 1439–1443.

Weltman, A. C., Bennett, N. M., Ackman, D. A., Misage, J. H., Campana, J. J., Fine, L. S., Doniger, A. S., Balzano, G. J. and Birkhead, G. S. 1996. An outbreak of hepatitis A associated with a bakery, New York, 1994: the 1968 'West Bransch, Michigan' outbreak repeated. Epidemiol. Infect., Vol. 117, pp. 333–341.

Whiting, R. C. and Buchanan, R. L. 1993. Letter to the editor: A classification of models in predictive microbiology – A reply to K. R. Davey. Food Microbiol., Vol. 10., pp. 175–177.

Wiedmann, M., Arvik, T., Bruce, J. L., Neubauer, J., del Piero, F., Smith, M. C., Hurley, J., Mohammed, H. O. and Batt, C. A. 1997. Investigation of a listeriosis epizootic in sheep in New York state. AJVR, Vol. 58, pp. 733–737.

Wirtanen, G. 1995. Biofilm formation and its elimination from food processing equipment. VTT Publications 251. Doctoral Thesis. Espoo: VTT. 106 p + app. 48 p. ISBN 951-38-4789-6.

Wirtanen, G. and Mattila-Sandholm, T. 1992a. Effect of the growth phase of foodborne biofilms on their resistance to a chlorine sanitizer. Part II. LWT, Vol. 25, pp. 50–54.

Wirtanen, G. and Mattila-Sandholm, T. 1992b. Removal of foodborne biofilms – Comparison of surface and suspension tests, part I. LWT, Vol. 25, pp. 43–49.

Wirtanen, G. and Mattila-Sandholm, T. 1993. Epifluorescence image analysis and cultivation of foodborne biofilm bacteria grown on stainless steel surfaces. J. Food Prot., Vol. 56, pp. 678–683.

Wirtanen, G. and Salo, S. (eds.). 2004. DairyNET – Hygiene control in Nordic dairies. VTT Publications 545. Espoo: VTT. 253 p. + app. 63 p. ISBN 951-38-6408-1. http://www.vtt.fi/inf/pdf/publications/2004/P545.pdf.

Worsfold, D. 2005. A survey of food safety training in small food manufacturers. Int. J. Environ. Health Res., Vol. 15, pp. 281–288.

Vorst, K. L., Todd, E. C. D. and Ryser, E. T. 2006a. Transfer of *Listeria monocytogenes* during mechanical slicing of turkey breast, bologna and salami. J. Food Prot., Vol. 69, pp. 619–626.

Vorst, K. L., Todd, E. C. D. and Ryser, E. T. 2006b. Transfer of *Listeria monocytogenes* during slicing of turkey breast, bologna and salami with simulated kitchen knives. J. Food Prot., Vol. 69, pp. 2939–2946.

WTO. 1994. Agreement on the application of sanutary and phytosanitary measures. Geneva, Switzerland: World Trade Organization.

Yano, K. 2005. Global listing program. New Food, Vol. 8, pp. 44-47.

Appendices of this publication are not included in the PDF version. Please order the printed version to get the complete publication (http://www.vtt.fi/publications/index.jsp)



Series title, number and report code of publication

VTT Publications 669 VTT-PUBS-669

Project number

Author(s) Aarnisalo, Kaarina

Title

Equipment hygiene and risk assessment measures as tools in the prevention of *Listeria monocytogenes* -contamination in food processes

Abstract

A problematic pathogen occurring in food processing is *Listeria monocytogenes*. Its efficient control at the processing plant level requires good equipment hygiene, including functioning good manufacturing and hygiene practices used by all employees, effective means of decontamination and rapid detection of contamination sources, as well as hazard analysis systems supported by risk assessment procedures. The present thesis focuses on deficiencies and improvements in these equipment hygiene and risk assessment practices with the aim of elucidating and developing the most efficient practices against *L. monocytogenes*.

The hygienically most problematic types of equipment in the Finnish food industry were identified as the packaging machines, conveyers, dispensers, slicing machines and cooling machines. The main reason for the equipment being considered as problematic was poor hygienic design. Equipment designers must focus their performance on more suitable equipment design. Clear deficiencies were also found in hygiene performance of the maintenance personnel in food processing plants. The results of these studies also indicate that L. monocytogenes may be transferred through maintenance work. Training of maintenance personnel with reference to hygienic practices must be increased. The results of the thesis showed that lubricants used in food processing equipment may act as contamination vehicles of L. monocytogenes. An investigation of the efficiency of eight commonly used commercial disinfectants against L. monocytogenes strains at +5 °C was performed. The tested agents were generally efficient at the recommended concentrations and effect times, with only a few exceptions. The suitability of automated ribotyping was also compared with the traditionally accepted and successfully used pulsed-field gel electrophoresis (PFGE) to discriminate L. monocytogenes isolates and thus trace contamination sources in food plants. PFGE had a higher discriminatory power for L. monocytogenes isolates than automated ribotyping. However, based on its automation and rapidity automated ribotyping can be considered a good method for control purposes. Additionally, risk assessment practices were developed by investigating and modelling recontamination of a product and by a plant-level quantitative risk assessment. Recontamination was modelled by investigating transfer of L. monocytogenes from slicing blade to slices of cold-salted salmon. Transfer with a progressive exponential reduction in the quantity of bacteria in slices was detected. As a concluding example, a practical approach to quantitative risk assessment of L. monocytogenes for one product at the plant level is presented. This approach helps food processors in illustrating the risks caused by the products for consumers.

ISBN

978-951-38-7069-0 (soft back ed.)

978-951-38-7070-6 (URL: http://www.vtt.fi/publications/index.jsp)

Series title and ISSN

VTT Publications 1235-0621 (soft back ed.) 1455-0849 (URL: http://www.vtt.fi/publications/index.jsp)

Date	Language	Pages			
December 2007 English, Finnish abstr.		101 p. + app. 65 p.			
Name of project		Commissioned by			
Keywords		Publisher			
Reywords					
food hygiene, Listeria mon	ocytogenes, risk	VTT Technical Research Centre of Finland			
food hygiene, <i>Listeria mon</i> assessment, hygienic desig	n, disinfectant,	VTT Technical Research Centre of Finland P.O. Box 1000, FI-02044 VTT, Finland			
food hygiene, <i>Listeria mon</i> assessment, hygienic desig lubricant, maintenance, mi	<i>ocytogenes</i> , risk n, disinfectant, crobial modelling,	VTT Technical Research Centre of Finland P.O. Box 1000, FI-02044 VTT, Finland Phone internat. +358 20 722 4520			



Tekijä(t)

Aarnisalo, Kaarina

Nimeke

Laitehygienia- ja riskinarviointitoimenpiteet työvälineinä *Listeria* monocytogenes -kontaminaation torjumisessa elintarvikeprosesseissa

Tiivistelmä

Listeria monocytogenes -bakteeri on elintarvikeprosesseissa esiintyvä ongelmallinen patogeeni. Sen tehokas hallinta tuotantolaitostasolla koostuu hyvästä laitehygieniasta ja sisältää toimivat, hyvät tuotanto- ja hygieniakäytännöt kaikkien työntekijöiden osalta, tehokkaat bakteerin tuhoamiseen tarkoitetut menetelmät ja nopeat kontaminaatiolähteiden toteamismenetelmät yhdistettynä riskinarviointikäytännöillä tuettuun prosessin vaara-analyysiin. Työssä selvitettiin puutteita ja parannuksia nykyisin käytössä olevien laitehygieniakäytäntöjen osalta ja kehitettiin riskinarviointikäytäntöjä mahdollisimman tehokkaiden *L. monocytogenes* -bakteerin torjumis-keinojen varmistamiseksi.

Tuotantohygienialtaan ongelmallisimmiksi laitteiksi Suomen elintarviketeollisuudessa todettiin pakkauskoneet, kuljettimet ja annostelu-, siivutus- ja jäähdytyskoneet. Pääasialliseksi syyksi todettiin huono hygieeninen laitesuunnittelu. Laitevalmistajien on kehitettävä elintarviketeollisuuteen tarkoitettujen laitteiden hygieenisyyttä. Työssä tehdyssä selvityksessä laitteita huoltavan kunnossapitohenkilöstön hygieenisistä toimintatavoista voitiin näissä tavoissa todeta selkeitä puutteita. Löydettiin myös viite siitä, että L. monocytogenes voisi siirtyä huoltotyön yhteydessä tehtaan sisällä. Kunnossapitohenkilöstön hygieniakoulutuksen määrää on lisättävä. Työn tulokset osoittivat myös, että tuotantolaitteistoissa käytettävät voiteluaineet voivat toimia L. monocytogenes -kontaminaation lähteinä ja siirtymisvälineenä. Työssä testatut käytössä olevat kaupalliset desinfiointiaineet (8 kpl) tehosivat L. monocytogenes -bakteeriin yleensä valmistajan antamilla alhaisimmilla käyttökonsentraatioilla ja käyttöajoilla +5 asteessa, vain joitakin poikkeuksia lukuun ottamatta. Automaattisen ribotyypityksen soveltuvuutta L. monocytogenes -isolaattien erotteluun ja siten kontaminaatiolähteiden jäljittämiseen elintarvikeprosesseissa verrattiin hyväksi todettuun pulssikenttägeelielektroforeesi (PFGE) -menetelmään, ja tulokset osoittivat, että PFGE oli erottelukyvyltään parempi. Nopeutensa ja automaattisuutensa takia automaattinen ribotyypitys soveltuu kuitenkin myös hyvin hygieniavalvontaan. Työssä kehitettiin lisäksi L. monocytogenes -bakteerin riskinarviointikäytäntöjä selvittämällä ja mallintamalla tuotteen jälkikontaminaatiota sekä kuvaamalla tehdastason riskinarviointi. Jälkikontaminaatiota mallinnettiin tutkimalla L. monocytogenes -bakteerin siirtymistä siivutettaessa terästä graavilohisiivuihin. Selkeää siirtymistä todettiin bakteerimäärien eksponentiaalisesti pienentyessä siivuissa. Viimeisenä esimerkkinä kuvataan käytännönläheinen yhden tuotteen L. monocytogenes -riskinarviointi. Tämä lähestymistapa auttaa elintarviketuottajia hahmottamaan kuluttajille tuotteista aiheutuvaa riskiä.

ISBN							
978-951-38-7069-0 (nid.)	978-951-38-7069-0 (nid.)						
978-951-38-7070-6 (URL: http://www.vtt.fi/publications/index.jsp)							
Avainnimeke ja ISSN			Projektinumero				
VTT Publications							
1235-0621 (nid.)							
1455-0849 (URL: http://w	ww.vtt.fi/publications/inc	lex.jsp)					
Julkaisuaika	Kieli	Sivuja					
Joulukuu 2007	Englanti, suom. tiiv.	101 s. + liitt. 65 s.					
Projektin nimi		Toimeksiantaja(t)					
Avainsanat		Julkaisija					
food hygiene, Listeria mor	<i>ocytogenes</i> , risk	VTT					
assessment, hygienic desig	n, disinfectant,	PL 1000, 02044 VTT					
lubricant, maintenance, mi	crobial modelling,	Puh. 020 722 4404					
transfer of bacteria, automa	ated ribotyping	Faksi 020 722 4374					

Julkaisun sarja, numero ja raporttikoodi

VTT Publications 669 VTT-PUBS-669

VTT PUBLICATIONS

- 650 Wessberg, Nina. Teollisuuden häiriöpäästöjen hallinnan kehittämishaasteet. 2007. 195 s. + liitt. 4 s.
- 651 Laitakari, Juhani. Dynamic context monitoring for adaptive and context-aware applications. 2007. 111 p. + app. 8 p.
- 652 Wilhelmson, Annika. The importance of oxygen availability in two plant-based bioprocesses: hairy root cultivation and malting. 2007. 66 p. + app. 56 p.
- 653 Ahlqvist, Toni, Carlsen, Henrik, Iversen, Jonas & Kristiansen, Ernst. Nordic ICT Foresight. Futures of the ICT environment and applications on the Nordic level. 2007. 147 p. + app. 24 p.
- 654 Arvas, Mikko. Comparative and functional genome analysis of fungi for development of the protein production host *Trichoderma reesei*. 100 p. + app. 105 p.
- 655 Kuisma, Veli Matti. Joustavan konepaja-automaation käyttöönoton onnistumisen edellytykset. 2007. 240 s. + liitt. 68 s.
- 656 Hybrid Media in Personal Management of Nutrition and Exercise. Report on the HyperFit Project. Ed. by Paula Järvinen. 121 p. + app. 2 p.
- 657 Szilvay, Géza R. Self-assembly of hydrophobin proteins from the fungus *Trichoderma reesei.* 2007. 64 p. + app. 43 p.
- 658 Palviainen, Marko. Technique for dynamic composition of content and contextsensitive mobile applications. Adaptive mobile browsers as a case study. 2007. 233 p.
- 659 Qu, Yang. System-level design and configuration management for run-time reconfigurable devices. 2007. 133 p.
- 660 Sihvonen, Markus. Adaptive personal service environment. 2007. 114 p. + app. 77 p.
- 661 Rautio, Jari. Development of rapid gene expression analysis and its application to bioprocess monitoring. 2007. 123 p. + app. 83 p.
- 662 Karjalainen, Sami. The characteristics of usable room temperature control. 2007. 133 p. + app. 71 p.
- Välkkynen, Pasi. Physical Selection in Ubiquitous Computing. 2007. 97 p. + app. 96 p.
- 664 Paaso, Janne. Moisture depth profiling in paper using near-infrared spectroscopy. 2007. 193 p. + app. 6 p.
- 666 Prunnila, Mika. Single and many-band effects in electron transport and energy relaxation in semiconductors. 2007. 68 p. + app. 49 p.
- 669 Aarnisalo, Kaarina. Equipment hygiene and risk assessment measures as tools in the prevention of *Listeria monocytogenes* -contamination in food processes. 2007. 101 p. + app. 65 p.
- 670 Kolari, Kai. Fabrication of silicon and glass devices for microfluidic bioanalytical applications. 2007. 100 s. + liitt. 72 s.

Julkaisu on saatavana	Publikationen distribueras av	This publication is available from
VTT	VTT	VTT
PL 1000	PB 1000	P.O. Box 1000
02044 VTT	02044 VTT	FI-02044 VTT, Finland
Puh. 020 722 4520	Tel. 020 722 4520	Phone internat. + 358 20 722 4520
http://www.vtt.fi	http://www.vtt.fi	http://www.vtt.fi

ISBN 978-951-38-7069-0 (soft back ed.) ISSN 1235-0621 (soft back ed.) ISBN 978-951-38-7070-6 (URL: http://www.vtt.fi/publications/index.jsp) ISSN 1455-0849 (URL: http://www.vtt.fi/publications/index.jsp)